

Dynamics of Rational Surface Automorphisms: Linear Fractional Recurrences

Eric Bedford* and Kyounghee Kim

§0. Introduction. Here we discuss automorphisms (biholomorphic maps) of compact, projective surfaces with positive entropy. Cantat [C1] has shown that the only (minimal dynamical system model) possibilities occur for tori and $K3$ (and certain of their quotients), and rational surfaces. $K3$ surfaces have been studied by Cantat [C2] and McMullen [M1]. Here we consider the family of birational maps of the plane which are defined by

$$f_{a,b} : (x, y) \mapsto \left(y, \frac{y+a}{x+b} \right), \quad (0.1)$$

and which provide an interesting source of automorphisms of rational surfaces. The maps $f_{a,b}$ form part of the family of so-called linear fractional recurrences, which were studied from the point of view of degree growth and periodicity in [BK]. We let $\mathcal{V} = \{(a, b) \in \mathbf{C}^2\}$ be the space of parameters for this family, and we define

$$\begin{aligned} q &= (-a, 0), \quad p = (-b, -a), \\ \mathcal{V}_n &= \{(a, b) \in \mathcal{V} : f_{a,b}^j q \neq p \text{ for } 0 \leq j < n, \text{ and } f_{a,b}^n q = p\}. \end{aligned} \quad (0.2)$$

In [BK] we showed that $f_{a,b}$ is birationally conjugate to an automorphism of a compact, complex surface $\mathcal{X}_{a,b}$ if and only if $(a, b) \in \mathcal{V}_n$ for some $n \geq 0$. The surface $\mathcal{X}_{a,b}$ is obtained by blowing up the projective plane $\mathbf{P}^2$ at the $n+3$ points $e_1 = [0 : 1 : 0]$, $e_2 = [0 : 0 : 1]$, and $f^j q$, $0 \leq j \leq n$. The dimension of $\text{Pic}(\mathcal{X}_{a,b})$ is thus $n+4$, and the action of $f_{a,b}^*$ on $\text{Pic}(\mathcal{X}_{a,b})$ is the same for all $(a, b) \in \mathcal{V}_n$: its characteristic polynomial is

$$\chi_n(x) := -1 + x^2 + x^3 - x^{1+n} - x^{2+n} + x^{4+n}. \quad (0.3)$$

This polynomial arises in the growth of Coxeter groups (see [F, p. 483]). When $n \geq 7$, χ_n has a root $\lambda_n > 1$, which is the unique root with modulus greater than one; and the entropy of $f_{a,b}$ is $\log \lambda_n > 0$. If $\psi_n(x)$ denotes the minimal polynomial of λ_n , then we may factor $\chi_n(x) = C_n(x)\psi_n(x)$, where C_n is a product of cyclotomic polynomials. The factorization of C_n into cyclotomic polynomials is given explicitly in Theorems 3.3 and 3.5.

McMullen [M2] noted that the maps $f_{a,b}$ provide representations of the Coxeter elements of certain Weyl groups, and he gave a construction of automorphisms which represent these Coxeter elements. By Theorem 8.7, the maps $f_{a,b}$ give essentially all possible rational surface automorphisms which represent these Coxeter elements.

The approach of this paper is to focus on the maps $f_{a,b}$ that have invariant curves. We obtain the following in Theorem 4.2:

* Research supported in part by the NSF.

Theorem A. *There are rational surface automorphisms $f_{a,b}$ which have positive entropy but have no invariant curves.*

We show in §4 that this gives a counterexample of a conjecture/question of Gizatullin, Harbourne, and McMullen.

We are also interested in the dynamical behavior of the maps with positive entropy. We say that a connected, invariant open set $\mathcal{D}$ is a rotation domain if the restriction $f|_{\mathcal{D}}$ is conjugate to a rotation of infinite order. (For mappings in dimension one, this corresponds to having a Siegel disk.) For a rotation domain $\mathcal{D}$, the closure of a generic orbit will be a smooth real torus $\mathbf{T}^d$ with d equal to 1 or 2, and we say that d is the rank of the rotation domain.

Theorem B. *Let $f_{a,b}$ be a mapping of the form (0.1) which is equivalent to an automorphism with positive entropy. If there is an $f_{a,b}$ -invariant curve, then exactly one of the following occurs:*

- (i) *f has a rank 1 rotation domain centered at one of the two fixed points.*
- (ii) *$a, b \in \mathbf{R}$, and the restriction f_R of f to the real points $\mathcal{X}_R$ has the same entropy as f . Further the (unique) invariant measure of maximal entropy is supported on a subset of $\mathcal{X}_R$ of zero area.*

As n increases, the maps with invariant cubics appear to form a very small fraction of the total number of maps in $\mathcal{V}_n$. It will be interesting to explore the dynamical properties of these other maps.

This paper is organized as follows. In §1 we show that if $S_{a,b}$ is a curve which is invariant under a map $f_{a,b}$ as in (0.1), then $S_{a,b}$ is cubic. We give the precise form of the cubic curve $S_{a,b}$ (if it exists) in §2. In §2, we also show that these parameter values (a, b) corresponding to invariant curves, correspond to $(a, b) \in \Gamma$, where Γ is the union of three specific curves in parameter space. In §3 we show that there is a close connection between the zeros of χ_n and the parameter values for which $f_{a,b}$ is an automorphism. In particular, we find the cyclotomic part C_n of χ_n . In §4 we show the existence of counterexamples to the Gizatullin/Harbourne/McMullen conjecture. And we prove Theorem 4.2, which also contains Theorem A above. We discuss the existence of rotation domains in §5. In §6 we show that the real maps with invariant curves have entropy equal to $\log \lambda_n$. In §7 we prove Theorem B. In §8 we discuss the connection with representations of the Coxeter element.

Acknowledgement. We thank Serge Cantat and Jeff Diller for explaining some of this material to us and giving helpful suggestions on this paper. We also thank the anonymous referees for their helpful comments.

§1. Invariant Curves. We will write our maps in projective coordinates $x = [x_0 : x_1 : x_2] = [1 : x : y]$. We define $\alpha = \gamma = (a, 0, 1)$ and $\beta = (b, 1, 0)$, so f is written

$$f_{a,b} : [x_0 : x_1 : x_2] \mapsto [x_0\beta \cdot x : x_2\beta \cdot x : x_0\alpha \cdot x].$$

The exceptional curves for the map f are given by the lines $\Sigma_0 = \{x_0 = 0\}$, $\Sigma_\beta = \{\beta \cdot x = 0\}$, and $\Sigma_\gamma = \{\gamma \cdot x = 0\}$. The indeterminacy locus $\mathcal{I}(f) = \{e_2, e_1, p\}$ consists of the vertices of the triangle $\Sigma_0\Sigma_\gamma\Sigma_\beta$. Let $\pi : \mathcal{Y} \rightarrow \mathbf{P}^2$ be the complex manifold obtained by

blowing up e_1 and e_2 , and let the exceptional fibers be denoted E_1 and E_2 . By Σ_0 , Σ_β and Σ_γ we denote the strict transforms in $\mathcal{Y}$. Let $f_{\mathcal{Y}} : \mathcal{Y} \rightarrow \mathcal{Y}$ be the induced birational map. Then the exceptional locus is Σ_γ , and the indeterminacy loci are $\mathcal{I}(f_{\mathcal{Y}}) = \{p\}$ and $\mathcal{I}(f_{\mathcal{Y}}^{-1}) = \{q\}$. In particular, $f_{\mathcal{Y}} : \Sigma_\beta \rightarrow E_2 \rightarrow \Sigma_0 \rightarrow E_1 \rightarrow \Sigma_B = \{x_2 = 0\}$. By curve, we mean an algebraic set of pure dimension 1, which may or may not be irreducible or connected. We say that an algebraic curve S is invariant if the closure of $f(S - \mathcal{I})$ is equal to S . We define the cubic polynomial $j_f := x_0(\beta \cdot x)(\gamma \cdot x)$, so $\{j_f = 0\}$ is the exceptional locus for f . For a homogeneous polynomial h we consider the condition that there exists $t \in \mathbf{C}^*$ such that

$$h \circ f = t \cdot j_f \cdot h. \quad (1.1)$$

Proposition 1.1. *Suppose that $(a, b) \notin \bigcup \mathcal{V}_n$, and S is an $f_{a,b}$ -invariant curve. Then S is a cubic containing e_1, e_2 , as well as $f^j q, f^{-j} p$ for all $j \geq 0$. Further, (1.1) holds for S .*

Proof. Let us pass to $f_{\mathcal{Y}}$, and let S denote its strict transform inside $\mathcal{Y}$. Since $(a, b) \notin \bigcup \mathcal{V}_n$, the backward orbit $\{f^{-n} p : n \geq 1\}$ is an infinite set which is disjoint from the indeterminacy locus of $\mathcal{I}(f_{\mathcal{Y}}^{-1})$. It follows that S cannot be singular at p (cf. Lemma 3.2 of [DJS]). Let μ denote the degree of S , and let $\mu_1 = S \cdot E_1$, $\mu_0 = S \cdot \Sigma_0$, and $\mu_2 = S \cdot E_2$. It follows that $\mu = \mu_2 + \mu_0 + \mu_1$. Further, since $f_{\mathcal{Y}} : E_2 \rightarrow \Sigma_0 \rightarrow E_1$, we must have $\mu_2 = \mu_0 = \mu_1$. Thus μ must be divisible by 3, and $\mu/3 = \mu_2 = \mu_0 = \mu_1$. Now, since S is nonsingular at $p = \Sigma_\beta \cap \Sigma_\gamma$, it must be transversal to either Σ_γ or Σ_β .

Let us suppose first that S is transversal to Σ_γ at p . Then we have $\mu = \mu_2 + S \cdot \Sigma_\gamma$. Thus S intersects $\Sigma_\gamma - \{p\}$ with multiplicity $\mu - (\mu/3) - 1$. If $\mu > 3$, then this number is at least 2. Now Σ_γ is exceptional, $f_{\mathcal{Y}}$ is regular on $\Sigma_\gamma - \{p\}$, and $f_{\mathcal{Y}}(\Sigma_\gamma - \{p\}) = q$. We conclude that S is singular at q . This is not possible by Lemma 3.2 of [DJS] since q is indeterminate for f^{-1} . This may also be seen because since $(a, b) \notin \bigcup \mathcal{V}_n$, it follows that $f^n q$ is an infinite orbit disjoint from the indeterminacy point p , which is a contradiction since S can have only finitely many singular points.

Finally, suppose that S is transversal to Σ_β . We have $\mu = S \cdot \Sigma_\beta + \mu_2$, and by transversality, this means that S intersects $\Sigma_\beta - \{p\}$ with multiplicity $\frac{2\mu}{3} - 1$. On the other hand, $f_{\mathcal{Y}}$ is regular on $\Sigma_\beta - \{p\}$, and $\Sigma_\beta - \{p\} \rightarrow E_2$. Thus the multiplicity of intersection of S with $\Sigma_\beta - \{p\}$ must equal the multiplicity of intersection with E_2 , but this is not consistent with the formulas unless $\mu = 3$. $\square$

The following was motivated by [DJS]:

Theorem 1.2. *Suppose that $(a, b) \in \mathcal{V}_n$ for some $n \geq 11$. If S is an invariant algebraic curve, then the degree of S is 3, and (1.1) holds.*

Proof. Let $\mathcal{X}$ be the manifold $\pi : \mathcal{X} \rightarrow \mathbf{P}^2$ obtained by blowing up $e_1, e_2, q, f q, \dots, f^n q = p$, and denote the blowup fibers by $E_1, E_2, Q, f Q, \dots, f^n Q = P$. Suppose that S is an invariant curve of degree m . By S , Σ_0 , etc., we denote the strict transforms of these curves inside $\mathcal{X}$. Let $f_{\mathcal{X}}$ be the induced automorphism of $\mathcal{X}$, so S is again invariant for $f_{\mathcal{X}}$, which we write again as f . Let us write the various intersection products with S as: $\mu_1 = S \cdot E_1$, $\mu_0 = S \cdot \Sigma_0$, $\mu_2 = S \cdot E_2$, $\mu_P = S \cdot P$, $\mu_\gamma = S \cdot \Sigma_\gamma$, $\mu_Q = S \cdot Q$. Since $e_1, e_2 \in \Sigma_0$, we have

$$\mu_1 + \mu_0 + \mu_2 = m.$$

Now we also have

$$f: \Sigma_\beta \rightarrow E_2 \rightarrow \Sigma_0 \rightarrow E_1$$

so $\mu_\beta = \mu_2 = \mu_0 = \mu_1 = \mu$ for some positive integer μ , and $m = 3\mu$. Similarly, $p, e_2 \in \Sigma_\beta$, so we conclude that $\mu_P + \mu_\beta + \mu_2 = m$, and thus $\mu_P = \mu$. Following the backward orbit of P , we deduce that $S \cdot f^j Q = \mu$ for all $0 \leq j \leq n$.

Now recall that if $L \in H^{1,1}(\mathbf{P}^2, \mathbf{Z})$ is the class of a line, then the canonical class of $\mathbf{P}^2$ is $-3L$. Thus the canonical class $K_{\mathcal{X}}$ of $\mathcal{X}$ is $-3L + \sum E$, where sum is taken over all blowup fibers E . In particular, the class of S in $H^{1,1}(\mathcal{X})$ is $-\mu K_{\mathcal{X}}$. Since we obtained $\mathcal{X}$ by performing $n + 3$ blowups on $\mathbf{P}^2$, the (arithmetic) genus formula, applied to the strict transform of S inside $\mathcal{X}$, gives:

$$g(S) = \frac{S \cdot (S + K_{\mathcal{X}})}{2} + 1 = \frac{\mu(\mu - 1)}{2} K_{\mathcal{X}}^2 + 1 = \frac{\mu(\mu - 1)}{2} (9 - (3 + n)) + 1.$$

Now let ν denote the number of connected components of S (the strict transform inside $\mathcal{X}$). We must have $g(S) \geq 1 - \nu$. Further, the degree 3μ of S must be at least as large as ν , which means that $\mu(\mu - 1)(n - 6) \leq 2\nu \leq 6\mu$ and therefore $\mu \leq 6/(n - 6) + 1$. We have two possibilities: (i) If $n \geq 13$, then $\mu = 1$, and S must have degree 3; (ii) if $n = 11$ or 12 , either $\mu = 1$, (i.e., the degree of S is 3), or $\mu = 2$. Let us suppose $n = 11$ or 12 and $\mu = 2$. From the genus formula we find that $5 \leq \nu \leq 6$. We treat these two cases separately.

Case 1. S cannot have 6 connected components. Suppose, to the contrary, $\nu = 6$. First we claim that S must be minimal, that is, we cannot have a nontrivial decomposition $S = S_1 \cup S_2$, where S_1 and S_2 are invariant. By the argument above, S_1 and S_2 must be cubics, and thus they must both contain all $n + 3 \geq 14$ points of blowup. But then they must have a common component, so S must be minimal.

Since the degree of S is 6, it follows that S is the union of 6 lines which map $L_1 \rightarrow L_2 \rightarrow \cdots \rightarrow L_6 \rightarrow L_1$. Further, each L_i must contain exactly one point of indeterminacy, since it maps forward to a line and not a quadric. Since the class of S in $H^{1,1}(\mathcal{X})$ is $-2K_{\mathcal{X}}$, we see that $e_1, e_2, p, q \in S$ with multiplicity 2. Without loss of generality, we may assume that $e_1 \in L_1$, which means $\Sigma_\beta \cap L_1 \neq \emptyset$, and therefore $e_2 \in L_2$. Similarly $q \in L_3$. Since the backward image of L_3 is a line, $e_1, e_2 \notin L_3$, and thus $p \in L_3$, which gives $\Sigma \cap L_3 \neq \emptyset$. Continuing this procedure, we end up with $L_6 \ni p, q$. It follows that $L_1 = L_4$, $L_2 = L_5$, and $L_3 = L_6$, so S has only 3 components.

Case 2. S cannot have 5 connected components. Suppose, to the contrary, that $\nu = 5$. It follows that S is a union of 4 lines and one quadric. Without loss of generality we may assume that $L_1 \rightarrow Q \rightarrow L_2 \rightarrow L_3$. Since L_1 maps to a quadric, it cannot contain a point of indeterminacy, which means that $L_1 \cap \Sigma_0 \neq \emptyset$, $L_1 \cap \Sigma_\beta \neq \emptyset$, and $L_1 \cap \Sigma_\gamma \neq \emptyset$. It follows that $e_1, e_2, q \in Q$. On the other hand, since L_2 maps to a quadric by f^{-1} , we have $e_1, e_2, p \in Q$, and $e_1, e_2, q \notin L_2$. Thus we have that $Q \cap \Sigma_0 = \{e_1, e_2\}$, $Q \cap \Sigma_\beta = \{e_2, p\}$, and $Q \cap \Sigma_\gamma = \{e_1, p\}$. It follows that $q \notin L_2$, which means that L_2 does not contain any point of indeterminacy, and therefore L_2 maps to a quadric.

Thus we conclude that S has degree 3, so we may write $S = \{h = 0\}$ for some cubic h . Since the class of S in $H^{1,1}(\mathcal{X})$ is $-3K_{\mathcal{X}}$, we see that $e_1, e_2, q \in S$. Since these are the images of the exceptional lines, the polynomial $h \circ f$ must vanish on $\Sigma_0 \cup \Sigma_\gamma \cup \Sigma_\beta$. Thus j_f divides $h \circ f$, and since $h \circ f$ has degree 6, we must have (1.1). $\square$

Remarks. (a) From the proof of Theorem 1.2, we see that if S is an invariant curve, $n \geq 11$, then S contains e_1 , e_2 , and $f^j q$, $0 \leq j \leq n$. (b) The only positive entropy parameters which are not covered in Theorem 1.2 are the cases $n = 7, 8, 9, 10$. By Proposition B.1, we have $\mathcal{V}_n \subset \Gamma$ for $7 \leq n \leq 10$.

Corollary 1.3. *If S is f -periodic with period k , and if $n \geq 11$, then $S \cup \dots \cup f^{k-1}S$ is invariant and thus a cubic.*

§2. Invariant Cubics. In this section, we identify the parameters $(a, b) \in \mathcal{V}$ for which the birational map $f_{a,b}$ has an invariant curve, and we look at the behavior of $f_{a,b}$ on this curve. We define the functions:

$$\begin{aligned} \varphi_1(t) &= \left(\frac{t - t^3 - t^4}{1 + 2t + t^2}, \frac{1 - t^5}{t^2 + t^3} \right), \\ \varphi_2(t) &= \left(\frac{t + t^2 + t^3}{1 + 2t + t^2}, \frac{-1 + t^3}{t + t^2} \right), \quad \varphi_3(t) = (1 + t, t - t^{-1}), \end{aligned} \quad (2.1)$$

The proofs of the results in this section involve some calculations that are possible but tedious to do by hand; but they are not hard with the help of Mathematica or Maple.

Theorem 2.1. *Let $t \neq 0, \pm 1$ with $t^3 \neq 1$ be given. Then there is a homogeneous cubic polynomial P satisfying (1.1) if and only if $(a, b) = \varphi_j(t)$ for some $1 \leq j \leq 3$. If this occurs, then (up to a constant multiple) P is given by (2.2) below.*

Proof. From the proof of Theorem 1.2, we know that P must vanish at e_1, e_2, q, p . Using the conditions $P(e_1) = P(e_2) = P(q) = 0$, we may set

$$\begin{aligned} P[x_0 : x_1 : x_2] &= (-a^2 C_1 + a C_2) x_0^3 + C_2 x_1 x_0^2 + C_3 x_2 x_0^2 + C_1 x_0 x_1^2 + \\ &\quad + C_4 x_2 x_1^2 + C_5 x_0 x_2^2 + C_6 x_1 x_2^2 + C_7 x_0 x_1 x_2 \end{aligned}$$

for some $C_1, \dots, C_7 \in \mathbb{C}$. Since $e_1, e_2, q \in \{P = 0\}$, we have $P \circ f = j_f \cdot \tilde{P}$ for some cubic $\tilde{P}$. A computation shows that

$$\begin{aligned} \tilde{P} &= (-ab^2 C_1 + b^2 C_2 + b C_3 + a C_5) x_0^3 + (-2ab C_1 + 2b C_2 + C_3) x_1 x_0^2 + \\ &\quad (b C_1 + C_5 + a C_6 + b C_7) x_2 x_0^2 + (-a C_1 + C_2) x_0 x_1^2 + C_1 x_2 x_1^2 + (b C_4 + C_6) x_0 x_2^2 + \\ &\quad + C_4 x_1 x_2^2 + (2b C_1 + C_7) x_0 x_1 x_2. \end{aligned}$$

Now setting $\tilde{P} = tP$ and comparing coefficients, we get a system of 8 linear equations in $C_1, \dots, C_7$ of the form

$$M \cdot [x_0^3, x_1 x_0^2, x_2 x_0^2, x_0 x_1^2, x_0 x_2^2, x_0 x_1 x_2, x_1 x_2^2, x_1^2 x_2]^t = 0.$$

We check that there exist cubic polynomials satisfying (1.1) if and only if the two principal minors of M vanish simultaneously, which means that

$$b(a + abt + abt^4 - b^2 t^4 - at^5 + bt^5) = 0$$

$$-1 + (1 - a - b)t + (a + b)t^2 + b^2t^3 + b^2t^4 + (a - 2b)t^5 + (1 - a + 2b)t^6 - t^7 = 0$$

Solving these two equations for a and b , we obtain φ_j , $j = 1, 2, 3$ as the only solutions, and then solving $M = 0$ we find that P must have the form:

$$\begin{aligned} P_{t,a,b}(x) = & ax_0^3(-1+t)t^4 + x_1x_2(-1+t)t(x_2+x_1t) \\ & + x_0[2bx_1x_2t^3 + x_1^2(-1+t)t^3 + x_2^2(-1+t)(1+bt)] \\ & + x_0^2(-1+t)t^3[a(x_1+x_2t) + t(x_1+(-2b+t)x_2)] \end{aligned} \quad (2.2)$$

which completes the proof. $\square$

Remark. Let us discuss the values of t omitted in Theorem 2.1. There is no nonzero solution to (1.1) if $t = 0$. If $t = 1$ or $t = -1$, then (1.1) is solvable iff $(a, b) \in \{b = 0\} \subset \mathcal{V}_0 \cup \mathcal{V}_1 \cup \mathcal{V}_6$. If $t = \omega$ is a primitive cube root of unity, then $(a, b) = (0, 0) = \varphi_2(\omega)$ or $(a, b) = (1 + \omega, \omega - \bar{\omega}) = \varphi_3(\omega)$. By [BK], $\varphi_3(\omega) \notin \mathcal{V}_0 \cup \dots \cup \mathcal{V}_6$, and by Theorem 3.1, $\varphi_3(\omega) \notin \bigcup_{n \geq 7} \mathcal{V}_n$.

If h satisfies (1.1), then we may define a meromorphic 2-form η_P on $\mathbf{P}^2$ by setting $\eta_h := \frac{dx \wedge dy}{h(1, x, y)}$ on the affine coordinate chart $[1 : x : y]$. Then η_h satisfies $t f^* \eta_h = \eta_h$. It follows that if the points $\{p_1, \dots, p_k\}$ form a k -cycle which is disjoint from $\{h = 0\}$, then the Jacobian determinant of f around this cycle will be t^{-k} .

Let $\Gamma_j = \{(a, b) = \varphi_j(t) : t \in \mathbf{C}\} \subset \mathcal{V}$ denote the curve corresponding to φ_j , and set $\Gamma := \Gamma_1 \cup \Gamma_2 \cup \Gamma_3$. Consistent with [DJS], we find that the cases Γ_j yield cubics with cusps, lines tangent to quadrics, and three lines passing through a point.




Figure 2.1. Orbit of q for family Γ_1 ; $1 < t < \delta_\star$.

Γ_1 : *Irreducible cubic with a cusp.* To discuss the family Γ_1 , let $(a, b) = \varphi_1(t)$ for some $t \in \mathbf{C}$. Then the fixed points of $f_{a,b}$ are $FP_s = (x_s, y_s)$, $x_s = y_s = t^3/(1+t)$ and $FP_r = (x_r, y_r)$, $x_r = y_r = (-1 + t^2 + t^3)/(t^2 + t^3)$. The eigenvalues of $Df_{a,b}(FP_s)$ are $\{t^2, t^3\}$. The invariant curve is $S = \{P_{t,a,b} = 0\}$, with P as in (2.2). This curve S contains FP_s and FP_r , and has a cusp at FP_s . The point q belongs to S , and thus the orbit $f^j q$ for all j until possibly we have $f^j q \in \mathcal{I}$. The 2-cycle and 3-cycle are disjoint from S , so the multipliers in (B.1) must satisfy $\mu_2^3 = \mu_3^2$, from which we determine that $\Gamma_1 \subset \mathcal{V}$ is a curve of degree 6.

We use the notation δ_* for the real root of $t^3 - t - 1$. Thus $1 \leq \lambda_n < \delta_*$, and the λ_n increase to δ_* as $n \rightarrow \infty$. The intersection of the cubic curve with $\mathbf{RP}^2$ is shown in Figure 2.1. The exceptional curves Σ_β and Σ_γ are used as axes, and we have chosen a modification of polar coordinates so that Σ_0 , the line at infinity, appears as the bounding circle of $\mathbf{RP}^2$. The points $FP_{s/r}, e_1, e_2, p, q, fq, f^2q, f^3q$ all belong to S , and Figure 2.1 gives their relative positions with respect to the triangle $\Sigma_\beta, \Sigma_\gamma, \Sigma_0$ for all $1 < t < \delta_*$. Since $t > 1$, the points $f^j q$ for $j \geq 4$ lie on the arc connecting $f^4 q$ and FP_r , and $f^j q$ approaches FP_r monotonically along this arc as $j \rightarrow \infty$. In case (a, b) belongs to $\mathcal{V}_n$, then $f^n q$ lands on p . The relative position of S_t with respect to the axes is stable for t in a large neighborhood of $[1, \delta_*]$. However, as t increases to δ_* , the fixed point FP_r moves down to p ; and for $t > \delta_*$, FP_r is in the third quadrant. And as t decreases to 1, f approaches the (integrable) map $(a, b) = (-1/4, 0) \in \mathcal{V}_6$. The family $\mathcal{V}_6$ will be discussed in Appendix A. When $0 < t < 1$, the point FP_s becomes attracting, and the relative positions of q and $f q$, etc., are reversed. Figure 2.1 will be useful in explaining the graph shown in Figure 6.1.

Γ_2 : *Line tangent to a quadric.* Next we suppose that $(a, b) = \varphi_2(t)$. We let $S = \{P_{t,a,b} = 0\}$ be the curve in (2.2). In this case, the curve is the union of a line $L = \{t^2 x_0 + t x_1 + x_2 = 0\}$ and a quadric Q . The fixed points are $FP_s = (x_s, y_s)$, $x_s = y_s = -t^2/(1+t)$ and $FP_r = (x_r, y_r)$, $x_r, y_r = (1+t+t^2)/(t+t^2)$. The eigenvalues of $Df_{a,b}$ at FP_s are $\{-t, -t^2\}$. The 3-cycle and FP_r are disjoint from S , so we have $\det(Df_{a,b} FP_r)^3 - \mu_3 = 0$ on Γ_2 , with μ_3 as in (B.1). Extracting an irreducible factor, we find that $\Gamma_2 \subset \mathcal{V}$ is a quartic.




Figure 2.2. Orbit of q for family Γ_2 ; $1 < t < \delta_*$.

Figure 2.2 gives for Γ_2 the information analogous to Figure 2.1. The principal difference with Figure 2.1 is that S contains an attracting 2-cycle; there is a segment σ inside the line connecting $f^3 q$ to one of the period-2 points, and there is an arc $\gamma \ni p$ inside the quadric connecting $f^4 q$ to the other period-2 point. Thus the points $f^{2j+1} q$ will approach the two-cycle monotonically inside σ as $j \rightarrow \infty$, and the points $f^{2j} q$ will approach the two-cycle monotonically inside γ . The picture of S with respect to the triangle $\Sigma_\beta, \Sigma_\gamma, \Sigma_0$ is stable for t in a large neighborhood of $[1, \delta_*]$. As t increases to δ_* , one of the points of the 2-cycle moves down to p . As t decreases to 1, q moves up (and $f^2 q$ moves down) to $e_2 \in \mathcal{I}$, and $f q$ moves down to Σ_0 . The case $t = 1$ is discussed in Appendix A.

$\psi(\zeta) = FP_s + \zeta R$. Now f^2 fixes FP_s and R , and the differential of f^2 has an eigenvalue t^2 in the eigenvector L , so we have $f^2\psi(\zeta) = \psi(t^2\zeta)$. Since $p, q \in Q$, we have $f q, f^{-1}p \in L$. We see that $\zeta_q := t^3/(1-t^2-t^3)$ and $\zeta_p := (t^3-t-1)^{-1}$ satisfy $\psi(\zeta_q) = f q$ and $\psi(\zeta_p) = f^{-1}p$. If $n = 2k$, then the condition $f^n q = f^{2k} q = p$ is equivalent to the condition $t^{2n-2}\zeta_q = \zeta_p$, which is equivalent to $\chi_n(t) = 0$.

Finally we consider the case $j = 1$ and set $(a, b) = \varphi_1(t)$. If we substitute these values of (a, b) into the formula (3.1), we obtain a polynomial $P(x)$ which is cubic in x and which has coefficients which are rational in t . In order to parametrize S by $\mathbf{C}$, we set $\psi(\zeta) = FP_s + \zeta A + \zeta^2 B + \zeta^3 FP_r$. We may solve for $A = A(t)$ and $B = B(t)$ such that $P(\psi(\zeta)) = 0$ for all ζ . Thus f fixes $\psi(0)$ and $\psi(\infty)$, and $f(\psi(\zeta)) = \psi(t\zeta)$. We set $\zeta_q := t^2/(1-t^2-t^3)$ and $\zeta_p := t/(t^3-t-1)$. The condition $f^n q = p$ is equivalent to $t^n \zeta_q = \zeta_p$, or $-t^{n+2}/(t^3+t^2-1) = t/(t^3-t-1)$, or $\chi_n(t) = 0$. $\square$

For each n , we let $\psi_n(t)$ denote the minimal polynomial of λ_n .

Theorem 3.2. *Let $t \neq 1$ be a root of χ_n for $n \geq 7$. Then either t is a root of ψ_n , or t is a root of χ_j for some $0 \leq j \leq 5$.*

Proof. Let t be a root of χ_n which is not a root of ψ_n . This root has modulus 1, so by Lemma 1.6 of [W], it must be a root of unity. We will show it is a root of χ_j for some $0 \leq j \leq 5$. First we note that $\chi_6(t) = (t-1)^3(t+1)(t^2+t+1)(t^4+t^3+t^2+t+1)$, and so every root of χ_6 is a root of χ_j for some $0 \leq j \leq 5$. Similarly, we may check that the Theorem is true for $7 \leq n \leq 13$. By induction, it suffices to show that if t is a root of unity, then it is a root of χ_j for some $0 \leq j \leq n-1$. By Theorem 3.1, we see that $\chi_n(t) = 0$ if and only if $t^n \zeta_q(t) = \zeta_p(t)$, where $\zeta_q(t) = t^2/(1-t^2-t^3)$ and $\zeta_p(t) = t/(t^3-t-1)$.

Claim 1: We may assume $t^n \neq \pm 1$. Otherwise, from $t^n \zeta_q(t) = \zeta_p(t)$ we have

$$t^2(t^3-t-1) \pm t(t^3+t^2-1) = 0.$$

In case we take “+”, the roots are also roots of χ_0 , and in case we take “-”, there are no roots of unity.

Claim 2: If $t^k = 1$ for some $0 \leq k \leq n-1$, then t is a root of χ_j for some $0 \leq k-1$. As in the proof of Theorem 2.2, the orbit of ζ_q is $\{\zeta_q, t\zeta_q, \dots, t^{k-1}\zeta_q\}$. Thus the condition that $t^k = 1$ means that $f^k q = p$, so $\chi_k(t) = 0$.

Claim 3: If t is a primitive k th root of unity, we cannot have $k > n$. Otherwise, we have $t^m \zeta_p = \zeta_q$ for $m = k-n$, which means that t is a root of $\xi_m := t^{m-1}(t^3+t^2-1) + t^3-t-1$. This means that t is a root of both polynomials:

$$\begin{aligned} \chi_n/(t-1) &= 1 + t - (t^3 + t^4 + \dots + t^n) + t^{n+2} + t^{n+3} \\ \xi_m/(t-1) &= 1 + 2t + 2t^2 + (t^3 + \dots + t^{n-2}) + 2t^{m-1} + 2t^m + t^{m+1}. \end{aligned}$$

We divide our argument into four sub-cases: $1 \leq m \leq 4$, $5 \leq m \leq n-2$, $n-1 \leq m \leq n+6$, and $n+6 < m$. If $1 \leq m \leq 4$, we may check directly that ξ_m does not have a k th root of unity with $k > n$. If $5 \leq m \leq n-2$, then $k = n+m \geq 2m+2$. We may assume that $t = e^{i\theta}$, with $\theta = 2\pi/k$. Thus $\Im m(t^j) > 0$ for $j = 1, \dots, m+1$. It follows that $\Im m(\xi_m(t)/(t-1)) > 0$, contradicting the assumption that t is a root.

If $n - 1 \leq m \leq n + 6$, we show that ξ_m does not vanish at a k -th root of unity by inspecting the remainder obtained after performing three steps of the Euclidean Algorithm on the polynomials χ_n and ξ_m .

This leaves the case $2n + 6 < k$, i.e. $m > n + 6$. Here we have

$$\begin{aligned} \xi_m/(t-1) + \chi_n/(t-1) &= 2 + 3t + 2t^2 + t^{n+1} + 2t^{n+2} + 2t^{n+3} + \\ &\quad + t^{n+4} + \dots + t^{m-4} + t^{m-2} + 2t^{m-1} + 2t^m + t^{m+1}. \end{aligned}$$

Since $t^k = 1$, and since $m = n - k$, we have $\Im m(t^{n+1} + t^{m-1}) = \Im m(t^{n+2} + t^{m-2}) = \dots = 0$. Applying this repeatedly, we find

$$\begin{aligned} \Im m(\xi_m/(t-1) + \chi_n/(t-1)) &= \Im m(3t + 2t^2 + t^{n+2} + t^{n+3} + t^{m-1} + 2t^m + t^{m+1}) \\ &= \Im m(3t + 2t^2 - t^{n-1} - 2t^n - t^{n+1} + t^{n+2} + t^{n+3}) = 0. \end{aligned}$$

Again taking $t = e^{i\theta}$ with $\theta = 2\pi/k$, the previous equation becomes

$$\begin{aligned} &3 \sin \theta + 2 \sin(2\theta) - \sin((n-1)\theta) - 2 \sin(n\theta) - \sin((n+1)\theta) + \\ &\quad + \sin((n+2)\theta) + \sin((n+3)\theta) \\ &= 3 \sin \theta + 2 \sin(2\theta) + 2 (\cos((n+1)\theta) + \cos((n+2)\theta)) \sin \theta \\ &\quad - \sin((n-1)\theta) - \sin(n\theta). \end{aligned}$$

By $2\theta/\pi < \sin \theta = \sin(\pi - \theta) < \theta$ for $0 < \theta < \pi/2$, this last expression is seen to be strictly negative since $\theta = 2\pi/k$ with $k > 2n + 6$ and $n \geq 13$. This completes the proof of Claim 3. $\square$

Remark. Let us recall the construction of surface automorphisms given in [M2]. It starts with a root t of ψ_n and a singular cubic, which is one of the three cases considered above: cusp cubic, line and quadric, or three lines. The automorphism is then uniquely determined by a certain marking of the cubic. Now if we choose $(a, b) = \psi_*(t)$, then the centers of blowup on the invariant curve in $\mathcal{X}_{a,b}$ correspond exactly to the marked points, and $f_{a,b}$ is equivalent to the resulting automorphism.

The following result gives the possibilities for the roots of $\chi_n(x)/\psi_n(x) \in \mathbf{Z}[x]$.

Theorem 3.3. *Let $t \neq 1$ be a root of χ_n with $n \geq 7$. Then t is either a root of ψ_n , or t is a root of some χ_j for $0 \leq j \leq 5$. Specifically, if t is not a root of ψ_n , then it is a k th root of unity corresponding to one of the following possibilities:*

- (i) $k = 2$, $t + 1 = 0$, in which case 2 divides n ;
- (ii) $k = 3$, $t^2 + t + 1 = 0$, in which case 3 divides n ;
- (iii) $k = 5$, $t^4 + t^3 + t^2 + t + 1 = 0$, in which case $n \equiv 1 \pmod{5}$;
- (iv) $k = 8$, $t^4 + 1 = 0$, in which case $n \equiv 2 \pmod{8}$;
- (v) $k = 12$, $t^4 - t^2 + 1 = 0$, in which case $n \equiv 3 \pmod{12}$;
- (vi) $k = 18$, $t^6 - t^3 + 1 = 0$, in which case $n \equiv 4 \pmod{18}$;
- (vii) $k = 30$, $t^8 + t^7 - t^5 - t^4 - t^3 + t + 1 = 0$, in which case $n \equiv 5 \pmod{30}$.

Conversely, for each $n \geq 7$ and k satisfying one of the conditions above, there is a corresponding root t of χ_n which is a k th root of unity.

Proof. Recall that $\chi_n(t) = 0$ if and only if $t^n \zeta_q(t) = \zeta_p(t)$ by Theorem 2.2. If t is a k -th root of unity, then $k < n$ and $\chi_j(t) = 0$ for some $0 \leq j \leq 5$. In case $j = 0$, $\zeta_p(t) = \zeta_q(t)$, and $(t+1)(t^2+t+1) = 0$. Thus $t^n \zeta_q(t) = \zeta_p(t)$ if and only if $t+1 = 0$ and 2 divides n , or $t^2+t+1 = 0$ and 3 divides n . Now let us write $k_j = 5, 8, 12, 18, 30$ for $j = 1, 2, 3, 4, 5$, respectively, in case we have $1 \leq j \leq 5$, $t^j \zeta_q(t) = \zeta_p(t)$, and $n \equiv j \pmod{k_j}$. Thus $t^n \zeta_q(t) = \zeta_p(t)$ if and only if $n \equiv j \pmod{k_j}$. That is, $t^n \zeta_q(t) = (t^{k_j})^n t^j \zeta_q(t) = t^j \zeta_q(t) = \zeta_p(t)$. $\square$

As a corollary, we see that the number of elements of $\Gamma_j \cap \mathcal{V}_n$ is determined by the number of Galois conjugates of λ_n .

Corollary 3.4. *If $n \geq 7$, and if $1 \leq j \leq 3$ divides n , then*

$$\Gamma_j \cap \mathcal{V}_n = \{\varphi_j(t) : t \text{ is a root of } \psi_n\}.$$

In particular, these sets are nonempty.

Theorem 3.5. *If $n \geq 7$, then every root of χ_n is simple. Thus the possibilities enumerated in Theorem 3.3 give the irreducible factorization of χ_n . In particular, since each of the factors in (i–vii) below can occur only once, we have $n - 26 \leq \deg(\psi_n) \leq n + 3$.*

Proof. If t is a root of χ_n , then either it is a root of ψ_n , which is irreducible, or it is one of the roots of unity listed in Theorem 3.3. We have

$$\chi'_n(t) = (n+4)t^{n+3} - (n+2)t^{n+1} - (n+1)t^n + 3t^2 + 2t.$$

Since $\chi'_n(1) = 6 - n$, 1 is a simple root. Now we check all the remaining cases:

- (i) 2 divides n : $t+1 = 0 \Rightarrow \chi'_n(t) = -2 - n \neq 0$.
- (ii) 3 divides n : $t^2+t+1 = 0 \Rightarrow \chi'_n(t) = 3t^2 - nt + 3 \neq 0$.
- (iii) $n \equiv 1 \pmod{5}$: $t^4+t^3+t^2+t+1 = 0 \Rightarrow \chi'_n(t) = (n+4)t^4 - (n-1)t^2 - (n-1)t \neq 0$.
- (iv) $n \equiv 2 \pmod{8}$: $t^4+1 = 0 \Rightarrow \chi'_n(t) = -(n+2)t^3 - (n-2)t^2 - (n+2)t \neq 0$.
- (v) $n \equiv 3 \pmod{12}$: $t^4-t^2+1 = 0 \Rightarrow \chi'_n(t) = -(n+1)t^3 - (n-1)t^2 + 2t - 2 \neq 0$.
- (vi) $n \equiv 4 \pmod{18}$: $t^6-t^3+1 = 0 \Rightarrow \chi'_n(t) = -(n+2)t^5 + 3t^4 + 3t^2 - (n+2)t \neq 0$.
- (vii) $n \equiv 5 \pmod{30}$: $t^8+t^7-t^5-t^4-t^3+t+1 = 0 \Rightarrow$
 $\chi'_n(t) = (n+4)t^8 - (n+2)t^6 - (n+1)t^5 + 3t^2 + 2t \neq 0$. $\square$

Example. The number $n = 26$ corresponds to cases (i), (iii), and (iv), so we see that $\chi_{26} = (t-1)(t+1)(t^4+1)(t^4+t^3+t^2+t+1)\psi_{26}$, so ψ_{26} has degree 20.

§4. Surfaces without Anti-PluriCanonical Section. A curve S is said to be a pluri-anticanonical curve if it is the zero set of a section of $\Gamma(\mathcal{X}, (-K_{\mathcal{X}})^{\otimes n})$ for some $n > 0$. We will say that $(\mathcal{X}, f)$ is minimal if whenever $\pi : \mathcal{X} \rightarrow \mathcal{X}'$ is a birational morphism mapping $(\mathcal{X}, f)$ to an automorphism $(\mathcal{X}', f')$, then π is an isomorphism. Gizatullin conjectured that if $\mathcal{X}$ is a rational surface which has an automorphism f such that f^* has infinite order on $\text{Pic}(\mathcal{X})$, then $\mathcal{X}$ should have an anti-canonical curve. Harbourne [H] gave a counterexample to this, but this counterexample is not minimal, and f has zero entropy. See [Zh] for positive results in the zero entropy case.

Proposition 4.1. *Let $\mathcal{X}$ be a rational surface with an automorphism f . Suppose that $\mathcal{X}$ admits a pluri-anticanonical section. Then there is an f -invariant curve.*

Proof. Suppose there is a pluri-anticanonical section. Then $\Gamma(\mathcal{X}, (-K_{\mathcal{X}})^{\otimes n})$ is a nontrivial finite dimensional vector space for some $n > 0$, and f induces a linear action on this space. Let η denote an eigenvalue of this action. Since $\mathcal{X}$ is a rational surface, $S = \{\eta = 0\}$ is a nontrivial curve, which must be invariant under f . $\square$

The following answers a question raised in [M2, §12].

Theorem 4.2. *There is a rational surface $\mathcal{X}$ and an automorphism f of $\mathcal{X}$ with positive entropy such that $(\mathcal{X}, f)$ is minimal, but there is no f -invariant curve. In particular, there is no pluri-canonical section.*

Proof. We start by finding a map without invariant curve. We consider $(a, b) \in \mathcal{V}_{11}$. Suppose that $(\mathcal{X}_{a,b}, f_{a,b})$ has an invariant curve S . Then by Theorem 1.2, S must be a cubic. By Theorem 3.1 we must have $(a, b) \in \Gamma_1$. That is, $(a, b) = \varphi_1(t)$ for some t . By Theorem 3.3, t is a root of the minimal polynomial ψ_{11} . By Theorem 3.5, $\psi_{11}(t) = \chi_{11}(t)/((t-1)(t^4+t^3+t^2+t+1))$ has degree 10, so $\mathcal{V}_{11} \cap \Gamma_1$ contains 10 elements. However, there are 12 elements in $\mathcal{V}_{11} - \Gamma_1$. Each of these gives an automorphism $(\mathcal{X}_{a,b}, f_{a,b})$ with entropy $\log \lambda_{11} > 0$ and with no invariant curve.

Next we claim that $(\mathcal{X}_{a,b}, f_{a,b})$ is minimal. Suppose that $\pi : \mathcal{X}_{a,b} \rightarrow \mathcal{Y}$ is a holomorphic map which is birational, and suppose that the induced map $f_{\mathcal{Y}}$ is an automorphism. Then there are finitely many points $P = \{p_1, \dots, p_N\}$ such that $\pi^{-1}p_j$ has positive dimension. It follows that P is invariant under $f_{\mathcal{Y}}$ and thus $\pi^{-1}P$ is an invariant curve for $f_{a,b}$. The nonexistence of invariant curves then shows that $(\mathcal{X}_{a,b}, f_{a,b})$ is minimal.

The nonexistence of an pluri-anticanonical section follows from Proposition 4.1. $\square$

Remark. By Proposition B.1 we cannot take $n \leq 10$ the proof of Theorem 4.2.

§5. Rotation Domains. Given an automorphism f of a compact surface $\mathcal{X}$, we define the Fatou set $\mathcal{F}$ to be the set of normality of the iterates $\{f^n : n \geq 0\}$. Let $\mathcal{D}$ be an invariant component of $\mathcal{F}$. We say that $\mathcal{D}$ is a rotation domain if $f_{\mathcal{D}}$ is conjugate to a linear rotation (cf. [BS1] and [FS], as well as Ueda [U], where this terminology is more completely justified). In this case, the normal limits of $f^n|_{\mathcal{D}}$ generate a compact abelian group. In our case, the map f does not have finite order, so the iterates generate a torus $\mathbf{T}^d$, with $d = 1$ or $d = 2$. We say that d is the rank of $\mathcal{D}$. The rank is equal to the dimension of the closure of a generic orbit of a point of $\mathcal{D}$. Let us start by proving the existence of rank 1 rotation domains.

Theorem 5.1. *Suppose that $n \geq 7$, j divides n , and $(a, b) \in \Gamma_j \cap \mathcal{V}_n$. That is, $(a, b) = \varphi_j(t)$ for some $t \in \mathbf{C}$. If $t \neq \lambda_n, \lambda_n^{-1}$ is a Galois conjugate of λ_n , then $f_{a,b}$ has a rotation domain of rank 1 centered at FP_s .*

Proof. There are three cases. We saw in §1 that the eigenvalues of $Df_{a,b}$ at FP_s are $\{t^2, t^3\}$ if $j = 1$; they are $\{-t, -t^2\}$ if $j = 2$ and $\{\omega t, \omega^2 t\}$ if $j = 3$. Since λ_n is a Salem number, the Galois conjugate t has modulus 1. Since t is not a root of unity, it satisfies the Diophantine condition

$$|1 - t^k| \geq C_0 k^{-\nu} \quad (5.1)$$

for some $C_0, \nu > 0$ and all $k \geq 2$. This is a classical result in number theory. A more recent proof (of a more general result) is given in Theorem 1 of [B]. We claim now that if η_1 and η_2 are the eigenvalues of $Df_{a,b}$ at FP_s , then for each $m = 1, 2$, we have

$$|\eta_m - \eta_1^{j_1} \eta_2^{j_2}| \geq C_0(j_1 + j_2)^{-\nu} \quad (5.2)$$

for some $C_0, \eta > 0$ and all $j_1, j_2 \in \mathbf{N}$ with $j_1 + j_2 \geq 2$. There are three cases to check: Γ_j , $j = 1, 2, 3$. In case $j = 1$, we have that $\eta_m - \eta_1^{j_1} \eta_2^{j_2}$ is equal to either $t^2 - t^{2j_1+3j_2} = t^2(1 - t^{2(j_1-1)+3j_2})$ or $t^3 - t^{2j_1+3j_2} = t^3(1 - t^{2j_1+3(j_2-1)})$. Since $j_1 + j_2 > 1$, we see that (5.2) is a consequence of (5.1). In the case $j = 2$, Df^2 has eigenvalues $\{t^2, t^4\}$, and in the case $j = 3$, Df^3 has eigenvalues $\{t^3, t^3\}$. In both of these cases we repeat the argument of the case $j = 1$. It then follows from Zehnder [Z2] that $f_{a,b}$ is holomorphically conjugate to the linear map $L = \text{diag}(\eta_1, \eta_2)$ in a neighborhood of FP_s . $\square$




Figure 5.1. Orbits of three points in the rank 1 rotation domain containing FP_s ; $(a, b) \in \mathcal{V}_7 \cap \Gamma_1$. Two projections.

For each $n \geq 7$ and each divisor $1 \leq j \leq 3$ of n , the only values of $(a, b) \in \mathcal{V}_n \cap \Gamma_j$ to which Theorem 5.1 does not apply are the two values $\varphi_j(\lambda_n)$ and $\varphi_j(\lambda_n^{-1})$. For all the other maps in $\mathcal{V}_n \cap \Gamma$, the Siegel domain $\mathcal{D} \ni FP_s$ is a component of both Fatou sets $\mathcal{F}(f)$ and $\mathcal{F}(f^{-1})$. For instance, if $j = 1$, then f is conjugate on $\mathcal{D}$ to the linear map $(z, w) \mapsto (t^2 z, t^3 w)$. Thus, in the linearizing coordinate, the orbit of a point of $\mathcal{D}$ will be dense in the curve $\{|z| = 1, w^2 = cz^3\}$, for some r and c . In particular, the closure of the orbit bounds an invariant (singular) complex disk. Three such orbits are shown in Figure 5.1.

Now let us discuss the other fixed point. Suppose that $(a, b) \in \Gamma_1 \cap \mathcal{V}_n$ and $\{\eta_1, \eta_2\}$ are the multipliers at FP_r . As was noted in the proof above, t satisfies (5.1), and so by Corollary B.5, both η_1 and η_2 satisfy (5.1). On the other hand, the resonance given by Theorem B.3 means that they do not satisfy (5.2), and thus we cannot conclude directly that f can be linearized in a neighborhood of FP_r . However, by Pöschel [P], there are holomorphic Siegel disks (of complex dimension one) $s_j : \{|\zeta| < r\} \rightarrow \mathcal{X}_{a,b}$, $j = 1, 2$, with the property that $s'_j(0)$ is the η_j eigenvector, and $f(s_j(\zeta)) = s_j(\eta_j \zeta)$. We note that one of these Siegel disks will lie in the invariant cubic itself and will thus intersect the rotation

domain about FP_s . By Theorem B.4 there are similar resonances between the multipliers for the 2- and 3-cycles, and thus similar Siegel disks, in the cases $\Gamma_2 \cap \mathcal{V}_n$ and $\Gamma_3 \cap \mathcal{V}_n$ respectively.

If $j = 2, 3$ and $(a, b) \in \Gamma_j \cap \mathcal{V}_n$, then at FP_r we have

$$Df_{a,b} = \begin{pmatrix} 0 & 1 \\ -\frac{1}{t} & \frac{1+t}{1+t+t^2} \end{pmatrix} \text{ for } j = 2, \text{ and } = \begin{pmatrix} 0 & 1 \\ -\frac{1}{t} & \frac{1}{t+1} \end{pmatrix} \text{ for } j = 3. \quad (5.3)$$

We let η_1, η_2 denote the eigenvalues of $Df_{a,b}$ at FP_r . For both $j = 2$ and $j = 3$ we have $\eta_1 \eta_2 = 1/t$.

Lemma 5.2. *Let $|t| = 1$, let $(a, b) = \varphi_j(t)$, and let η_1, η_2 be the eigenvalues of $Df_{a,b}$ at FP_r .*

1. *If $j = 2$, then: $|\eta_1| = |\eta_2| = 1 \Leftrightarrow \operatorname{Re}(t) \geq -7/8$.*
2. *If $j = 3$, then: $|\eta_1| = |\eta_2| = 1 \Leftrightarrow \operatorname{Re}(t) \notin (-\frac{23-\sqrt{17}}{32}, \frac{-23+\sqrt{17}}{32})$.*

Proof. Set $t = e^{i\theta}$. Since $\eta_1 \eta_2 = 1/t$, we may set $\eta_1 = \delta e^{i\omega}$, $\eta_2 = \frac{1}{\delta} e^{-i(\theta+\omega)}$, $\delta > 0$. Consider first the case $j = 3$. With η_1, η_2 written as above, it follows from (5.3) that they are the eigenvalues if and only if $\eta_1 + \eta_2 = 1/(t+1)$. Setting $\delta = 1$, then, we have

$$\begin{aligned} \eta_1 + \eta_2 = \frac{1}{t+1} &\Leftrightarrow -1 + 2\cos\omega + 2\cos(\theta + \omega) = 0 \\ &\Leftrightarrow \cos\omega = \frac{1}{4} \pm \frac{\sqrt{-(\cos^2\theta - 1)(8\cos\theta + 7)}}{4(1 + \cos\theta)}. \end{aligned}$$

Thus there are solutions for ω if and only if the right hand side of the last equation is between -1 and 1 . We may check that this happens if and only if $\cos\theta \geq -7/8$, which gives condition (2).

The case $j = 2$ is similar. Again we set $\delta = 1$, and this time we find ω so that we have

$$\begin{aligned} \eta_1 + \eta_2 = \frac{1+t}{1+t+t^2} &\Leftrightarrow -\cos\frac{\theta}{2} + (1 + 2\cos\theta)\cos(\frac{\theta}{2} + \omega) = 0 \\ &\Leftrightarrow \cos\omega = \frac{\tau^2 \pm \sqrt{(1-\tau^2)(4\tau^2 + \tau - 1)(4\tau^2 - \tau - 1)}}{4\tau^2 - 1}, \end{aligned}$$

where we set $\tau = \cos\frac{\theta}{2}$ in the last equation. Solving for the right hand side of this last equation to be between -1 and 1 , we find condition (1). $\square$

Theorem 5.3. *Suppose that $n \geq 8$, $j = 2$ or 3 , and j divides n . Suppose that $(a, b) = \varphi_j(t)$, and $|t| = 1$ is a root of ψ_n which satisfies the condition of Lemma 5.2. Then $f_{a,b}$ has a rotation domain of rank 2 centered at FP_r .*

Proof. By Lemma 5.2, the multipliers η_1, η_2 of $Df_{a,b}$ at FP_r both have modulus 1. We will show that $\eta_1^{m_1} \eta_2^{m_2} \neq 1$ for all $m_1, m_2 \in \mathbf{Z}$, not both zero. Thus η_1 and η_2 are multiplicatively independent, and the rank of a rotation domain with such multipliers must be 2. In particular, we then have the non-resonance condition, which asserts that for $m = 1, 2$, we have $\eta_m - \eta_1^{j_1} \eta_2^{j_2} \neq 0$ for all $j_1, j_2 \in \mathbf{N}$ with $j_1 + j_2 \geq 2$. Since η_1 and

η_2 are algebraic, it follows from Theorem 1 of [B] that (5.2) holds. Thus by [Z2], $f_{a,b}$ is linearizable at FP_τ .

To show multiplicative independence, let us start with the case $j = 3$. Suppose that $\eta_1^{m_1} \eta_2^{m_2} = 1$ with $m_1 \geq m_2$ not both zero. We may rewrite this as

$$\eta_1^{m_1-m_2} (\eta_1 \eta_2)^{m_2} = 1 = \eta_1^{m_1-m_2} t^{-m_2}.$$

Choose τ such that $\tau^{m_1-m_2} = t$. Thus $\tau^{m_2} = \eta_1$ and $\tau^{-m_1} = \eta_2$. It follows that τ is a root of the following equations:

$$\psi_n(x^{m_1-m_2}) = 0, \text{ and } x^{m_2} + x^{-m_1} = \frac{1}{1 + x^{m_1-m_2}} \quad (5.4)$$

We know that $\psi_n(x^{m_1-m_2})$ has exactly $m_1 - m_2$ roots outside the closed unit disk, and these are $\lambda_n^{\frac{1}{m_1-m_2}}, \lambda_n^{\frac{1}{m_1-m_2}} \omega, \dots, \lambda_n^{\frac{1}{m_1-m_2}} \omega^{m_1-m_2-1}$, where $\omega^{m_1-m_2} = 1$.

Now let $\psi_n(x^{m_1-m_2}) = \phi_1(x) \cdots \phi_s(x)$, $\phi_j \in \mathbf{Z}[x]$ be the irreducible factorization. We may assume that τ is a root of ϕ_1 . Since τ is not a root of unity, the roots of ϕ_1 cannot all lie on the unit circle, and thus ϕ_1 must have a root $\tau_* = \lambda_n^{\frac{1}{m_1-m_2}} \omega^\ell$ for some ℓ . Since τ satisfies the second equation in (5.4), and ϕ_1 is irreducible, it follows that τ_* also satisfies this equation, which may be rewritten as

$$\tau_*^{m_2} + \frac{1}{\lambda_n} \frac{1}{\tau_*^{m_2}} = \frac{1}{1 + \lambda_n}.$$

Now we write $\tau_*^{m_2} = \delta e^{i\theta}$ with $\delta > 0$, $\delta \neq 1$, so this equation becomes

$$\delta(\cos \theta + i \sin \theta) + \frac{1}{\lambda_n \delta}(\cos \theta - i \sin \theta) = \frac{1}{1 + \lambda_n}.$$

The imaginary part is $\delta \sin \theta - \frac{1}{\lambda_n \delta} \sin \theta = 0$, so we must have either $\sin \theta = 0$ or $\delta = \lambda_n^{-1/2}$.

The first case is impossible. In the second case we have $\lambda_n^{-1/2} = \delta = \lambda_n^{m_2/(m_1-m_2)}$, which means that $m_1 = -m_2$. By our condition of multiplicative dependence, we find that $\eta_1 = \eta_2$, which implies that $4t^2 + 7t + 4 = 0$, which is impossible.

The case $j = 2$ is similar. Suppose again that we have a multiplicative dependence $\eta_1^{m_1} \eta_2^{m_2} = 1$, and we choose τ as before. Then τ is a root of

$$\psi_n(x^{m_1-m_2}) = 0, \text{ and } x^{m_2} + x^{-m_1} = \frac{1 + x^{m_1-m_2}}{1 + x^{m_1-m_2} + x^{2(m_1-m_2)}}.$$

We now have τ_* as before, which must satisfy the second equation, which gives

$$\tau_*^{m_2} + \frac{1}{\lambda_n} \tau_*^{-m_2} = \frac{1 + \lambda_n}{1 + \lambda_n + \lambda_n^2}.$$

Again, we set $\tau_*^{m_2} = \delta e^{i\theta}$, so this equation becomes

$$\delta(\cos \theta + i \sin \theta) + \frac{1}{\lambda_n \delta}(\cos \theta - i \sin \theta) = \frac{1 + \lambda_n}{1 + \lambda_n + \lambda_n^2}.$$

Setting the imaginary part equal to zero gives $\sin \theta = 0$ or $\delta = \lambda_n^{-1/2}$. The first possibility is impossible. The second possibility leads to the conclusion that $\eta_1 = \eta_2$, which means that $4t^4 + 7t^3 + 10t^2 + 7t + 4 = 0$. This is impossible, so η_1 and η_2 are multiplicatively independent. $\square$

Lemma 5.4. *Suppose that $n \geq 8$, and let $j = 2$ or 3 . If j divides n , then there exist $(a, b) \in \Gamma_j \cap \mathcal{V}_n$ such that $|\eta_1| = |\eta_2| = 1$.*

Proof. Let us recall the factorization $\chi_n = \tilde{C}_n \psi_n$, with $\tilde{C}_n$ given as in Theorem 3.5. Thus the roots of χ_n are given by $\lambda_n, \lambda_n^{-1}, 1$, the roots of unity $\{r_1, \dots, r_N\}$ coming from $\tilde{C}_n$, and those roots $\{t_1, \dots, t_M\}$ of ψ_n which have modulus 1. Since the sum of the roots of χ_n is zero, we must have

$$\lambda_n + \lambda_n^{-1} + 1 + \operatorname{Re} \sum r_\ell + \operatorname{Re} \sum t_\ell = 0.$$

On the other hand, $1 < \lambda_n < 1.4$, and the roots of each cyclotomic factor in $\tilde{C}_n$ add to either 0 or -1 . Thus we have

$$3.4 + \operatorname{Re} \sum t_\ell \geq 0.$$

Now we claim that we must have a value t_ℓ which satisfies both conditions (1) and (2) in Lemma 2.2. Otherwise, we have $\operatorname{Re}(t_\ell) \leq -5.8$ for all ℓ , which means that $M \leq 3.4/.58$, so that $M \leq 6$. By Theorem 3.5, the degree of ψ_n is at least $n - 26$, so we must have $n \leq 34$. The cases $8 \leq n \leq 34$ may be checked directly. $\square$

The following result is a combination of Theorems 5.1, 5.3 and Lemma 5.4; Theorem 10.1 of [M2] gives rank 2 rotation domains under the additional assumption that n is sufficiently large.

Theorem 5.5. *Suppose that $n \geq 8$, $j = 2$ or 3 , and j divides n . Then there exists $(a, b) \in \Gamma_j \cap \mathcal{V}_n$ such that $f_{a,b}$ has a rank 2 rotation domain centered at FP_r , as well as a rank 1 rotation domain centered at FP_s .*

§6. Real Mappings of Maximal Entropy. Here we consider real parameters $(a, b) \in \mathbf{R}^2 \cap \mathcal{V}_n$ for $n \geq 7$. Given such (a, b) , we let $\mathcal{X}_R$ denote the closure of $\mathbf{R}^2$ inside $\mathcal{X}_{a,b}$. We let $\lambda_n > 1$ be the largest root of χ_n , and for $1 \leq j \leq 3$, we let $f_{j,R}$ denote the automorphism of $\mathcal{X}_R$ obtained by restricting $f_{a,b}$ to $\mathcal{X}_R$, with $(a, b) = \varphi_j(\lambda_n)$.

Theorem 6.1. *There is a homology class $\eta \in H_1(\mathcal{X}_R)$ such that $f_{1,R*}\eta = -\lambda_n\eta$. In particular, $f_{1,R}$ has entropy $\log \lambda_n$.*

Proof. We use an octagon in Figure 6.1 to represent $\mathcal{X}_R$. Namely, we start with the real projective plane $\mathbf{RP}^2$; we identify antipodal points in the four “slanted” sides. The horizontal and vertical pairs of sides of the octagon represent the blowup fibers over the points e_1 and e_2 . These are labeled E_1 and E_2 ; the letters along the boundary indicate the identifications. (Since we are in a blowup fiber, the identification is no longer “antipodal.”) Further, the points $f^j q$ (written “ j ”) $0 \leq j \leq n$, are blown up, although we do not draw the blowup fibers explicitly. To see the relative positions of “ j ” with respect to the triangle $\Sigma_\beta, \Sigma_\gamma, \Sigma_0$, consult Figure 2.1. The 1-chains of the homology class η are represented by the directed graph $\mathcal{G}_1$ inside the manifold $\mathcal{X}_R$. If we project $\mathcal{G}_1 \subset \mathcal{X}_R$ down to the projective plane, then all of the incoming arrows at a center of blowup “ j ” will be tangent to each

other, as well as the outgoing arrows. Thus the projection of $\mathcal{G}_1$ will look like a train track.




Figure 6.1. Graph $\mathcal{G}_1$; Invariant homology class for family Γ_1 .

In order to specify the homology class η , we need to assign real weights to each edge of the graph. By “51” we denote the edge connecting “5” and “1”; and “170 = 1a70” denotes the segment starting at “1”, passing through “a”, continuing through “7”, and ending at “0”. Abusing notation, also write “51”, etc., to denote the weight of the edge, as well as the edge itself. We determine the weights by mapping η forward. We find that, upon mapping by f , the orientations of all arcs are reversed. Let us describe how to do this. Consider the arc “34” = “3e4”. The point “e” belongs to Σ_0 , and so it maps to E_1 . Thus “34” is mapped to something starting at “4”, passing through E_1 , and then continuing to “5”. Thus we see that “34” is mapped (up to homotopy) to “4f05”. Thus, the image of “34” covers “04” and “05”.

Inspection shows that no other arc maps across “04”, so we write “04 $\rightarrow$ 34” to indicate that the weight of side “04” in $f_*\eta$ is equal to the weight of “34”. Inspecting the images of all the arcs, we find that “24” also maps across “05”, so we write “05 $\rightarrow$ 24 + 34” to indicate how the weights transform as we push $\mathcal{G}_1$ forward. Looking at all possible arcs, we write the transformation $\eta \mapsto f_*\eta$ as follows:

$$\begin{aligned}
02 &\rightarrow 16 + 170 + \cdots + 1(n-1)0, & 03 &\rightarrow 24 + 25 + 26, & 04 &\rightarrow 34, \\
05 &\rightarrow 24 + 34, & 06 &\rightarrow 25, & 12 &\rightarrow 1n0, & 13 &\rightarrow 02, & 14 &\rightarrow 03, \\
15 &\rightarrow 04, & 16 &\rightarrow 05, & 170 &\rightarrow 06, & 1k0 &\rightarrow 1(k-1)0, & 7 < k &\leq n, \\
23 &\rightarrow 12, & 24 &\rightarrow 13, & 25 &\rightarrow 14, & 26 &\rightarrow 15, & 34 &\rightarrow 23,
\end{aligned} \tag{6.1}$$

The formula (6.1) defines a linear transformation on the space of coefficients of the 1-chains defining η . The spectral radius of the transformation (6.1) is computed in Appendix C, where we find that it is λ_n . Now let w denote the eigenvector of weights corresponding to the eigenvalue λ_n . It follows that if we assign these weights to η , then by construction we have $f_{1,R*}\eta = -\lambda_n\eta$, and η is closed. $\square$

Remark. Let us compare with the situation for real Hénon maps. In [BLS] it was shown that a real Hénon map has maximal entropy if and only if all periodic points are real. On the other hand, if $(a, b) = \varphi_1(t)$, $1 \leq t \leq 2$, the (unique) 2-cycle of the map $f_{a,b}$ is non-real. This includes all the maps discussed in Theorem 6.1, since all values of $t = \lambda_n$ are in this interval.




Figure 6.2. Invariant graph $\mathcal{G}_2$: $n = 2k$.

Theorem 6.2. The maps $f_{2,R}$ (if n is even) and $f_{3,R}$ (if n is divisible by 3) have entropy $\log \lambda_n$.

Proof. Since the entropy of the complex map f on $\mathcal{X}$ is $\log \lambda_n$, the entropies of $f_{2,R}$ and $f_{3,R}$ are bounded above by $\log \lambda_n$. In order to show that equality holds for the entropy of the real maps, it suffices by Yomdin's Theorem [Y] (see also [G]) to show that $f_{j,R}$ expands lengths by an asymptotic factor of λ_n . We will do this by producing graphs $\mathcal{G}_2$ and $\mathcal{G}_3$ on which f has this expansion factor. We start with the case $n = 2k$; the graph $\mathcal{G}_2$ is shown in Figure 6.2, which should be compared with Figure 2.2. We use the notation $01 = 0d1$ for the edge in $\mathcal{G}_2$ connecting “0” to “1” by passing through d . In this case, the notation already defines the edge uniquely; we have added the d by way of explanation. We will use $\mathcal{G}_2$ to measure length growth; the edges of $\mathcal{G}_2$ will be analogous to the “u-arcs” that were used in [BD2]. By “01” we will mean all arcs in $\mathcal{X}_R$ which connect the fiber “0” to the fiber “1”, and whose projections to $\mathbf{RP}^2$ are homotopic to the edge 01. In order to measure length growth, we will start with a configuration of arcs corresponding to edges of the graph. The symbol “01” will also be used in formulas (6.2) and (6.3), for instance, to denote the number of arcs which are of type 01.

Now we discuss how these arcs are mapped. The arc 01 crosses Σ_β and then $E_2 \ni d$ before continuing to “1”. Since Σ_β is mapped to E_2 and E_2 is mapped to Σ_0 , the image of 01 will start at “1” and cross E_2 and then Σ_0 before reaching “2”. Up to homotopy, we may slide the intersection points in E_2 and Σ_0 over to a point $g \in E_2 \cap \Sigma_0$. Thus, up to homotopy, f maps the edge 01 in $\mathcal{G}_2$ to the edge $1g2$.




Figure 6.3. Invariant graph $\mathcal{G}_3$: $n = 3k$.

Similarly, we see that the arcs $04, 06, \dots, 0(2k-1)$ all cross Σ_β and then Σ_γ . Thus the images of all these arcs will start at 1, pass through E_2 at d , then 0, and continue to the respective endpoints $5, 7, \dots, (2k-1)$. Since the images of all these arcs, up to homotopy, contain the edge 01 , the transformation of weights in the graph is given by the first entry of (6.2), and the whole transformation is given by the rest of (6.2):

$$\begin{aligned}
 01 &\rightarrow 04 + 061 + 081 + \dots + 0(2k-1)1, & 03 &\rightarrow 25 + 072 + 092 + \dots + 0(2k-1)2 \\
 04 &\rightarrow 3b4, & 061 &\rightarrow 25 + 45, & 12 &\rightarrow 0(2k)1, & 1c2 &\rightarrow 01, & 14 &\rightarrow 03 \\
 0(2j)1 &\rightarrow 0(2j-1)2, \quad j = 4, 5, \dots, k, & 0(2j+1)2 &\rightarrow 0(2j)1, \quad j = 3, 4, \dots, k-1 \\
 2a3 &\rightarrow 12, & 2e3 &\rightarrow 1c2, & 25 &\rightarrow 14, & 3b4 &\rightarrow 2e3, & 34 &\rightarrow 2a3, & 45 &\rightarrow 34.
 \end{aligned} \tag{6.2}$$

The characteristic polynomial for the transformation defined in (6.2) is computed in the Appendix, and the largest eigenvalue of (6.2) is λ_n , so $f_{2,R}$ has the desired expansion.

The case $n = 3k$ is similar. The graph $\mathcal{G}_3$ is given in Figure 6.3. Up to homotopy, $f_{3,R}$ maps the graph $\mathcal{G}_3$ to itself according to:

$$\begin{aligned}
 01 &\rightarrow 0b4, & 02 &\rightarrow 13 + 162 + 192 + \dots + 1(3k-3)2, \\
 0b4 &\rightarrow 3a4 + 073 + 0(10)3 + \dots + 0(3k-2)3, & 051 &\rightarrow 0b4 + 3c4 + 3a4, \\
 0(3j-1)1 &\rightarrow 0(3j-2)3, \quad j = 3, 4, \dots, k, & 1(3j)2 &\rightarrow 0(3j-1)1, \quad j = 2, 3, \dots, k, \\
 0(3j+1)3 &\rightarrow 1(3j)2, \quad j = 2, 3, \dots, k-1, \\
 12 &\rightarrow 01, & 13 &\rightarrow 02, & 23 &\rightarrow 1(3k)2, & 2d3 &\rightarrow 12, & 3a4 &\rightarrow 23, & 3c4 &\rightarrow 2d3.
 \end{aligned} \tag{6.3}$$

The linear transformation corresponding to (6.3) is shown in Appendix C to have spectral radius equal to λ_n , so $f_{3,R}$ has entropy $\log \lambda_n$. $\square$

§7. Proof of Theorem B. This Theorem is a consequence of results we have proved already. Let $f_{a,b}$ be of the form (0.1). By Proposition B.1 in Appendix B, we may suppose that $n \geq 11$. Thus if f has an invariant curve, then by Theorem 1.2, it has an invariant cubic, which is given explicitly by Theorem 2.1. Further, by Theorem 2.2, we must have $(a, b) = \varphi_j(t)$ for some j dividing n , and a value $t \in \mathbf{C}$ which is a root of χ_n . By Theorem 3.3, t cannot be a root of unity. Thus it is a Galois conjugate of λ_n . The Galois conjugates of λ_n are of two forms: either t is equal to λ_n or λ_n^{-1} , or t has modulus equal to 1.

In the first case, $(a, b) \in \mathcal{V} \cap \mathbf{R}^2$, and thus f is a real mapping. The three possibilities are $(a, b) \in \mathcal{V}_n \cap \Gamma_j$, $j = 1, 2, 3$, and these are treated in §5. In all cases, we find that the entropy of the real mapping $f_{R,j}$ has entropy equal to $\log \lambda_n$. By Cantat [C2], there is a unique measure μ of maximal entropy for the complex mapping. Since $f_{R,j}$ has a measure ν of entropy $\log \lambda_n$, it follows that $\mu = \nu$, and thus μ is supported on the real points. On the other hand, we know that μ is disjoint from the Fatou sets of f and f^{-1} . McMullen [M2] has shown that the complement of one of the Fatou sets $\mathcal{F}(f)$ or $\mathcal{F}(f^{-1})$ has zero volume. The same argument shows that the complement inside $\mathbf{R}^2$ has zero area. Thus the support of μ has zero planar area.

The other possibility is that t has modulus 1. In this case, Theorem B is a consequence of Theorem 5.1. $\square$

§8. Representing the Coxeter Element. Now we suppose that $\mathcal{X}$ is a rational surface, and $F \in \text{Aut}(\mathcal{X})$ is an automorphism such that the induced map F_* on $\text{Pic}(\mathcal{X})$ has infinite order. Then by a Theorem of Nagata there is a biholomorphic morphism $\pi : \mathcal{X} \rightarrow \mathbf{P}^2$ such that π can be factored $\pi_1 \circ \cdots \circ \pi_N$ into a composition of point blow-ups $\pi_\sigma : \mathcal{X}_\sigma \rightarrow \mathcal{X}_{\sigma-1}$ such that $\mathcal{X}_N = \mathcal{X}$ and $\mathcal{X}_0 = \mathbf{P}^2$. For each $1 \leq \sigma \leq N$ we let E_σ denote the exceptional blowup fiber for π_σ . If we let E_0 denote the class of a general line, then $\{E_\sigma : 0 \leq \sigma \leq N\}$ is a basis of $\text{Pic}(\mathcal{X})$. We may also define $\mathcal{E}_\sigma := (\pi_{\sigma+1} \circ \cdots \circ \pi_N)^* E_\sigma$, and $\mathcal{E}_0 = E_0$. Thus $\{\mathcal{E}_\sigma : 0 \leq \sigma \leq N\}$ is a geometric basis of $\text{Pic}(\mathcal{X})$, which means that $\mathcal{E}_0^2 = 1$, $\mathcal{E}_\sigma^2 = -1$ for $1 \leq \sigma \leq N$, and $\mathcal{E}_i \cdot \mathcal{E}_j = 0$ when $i \neq j$.

Another result of Nagata says that F_* must belong to the Weyl group W_N , which is generated by reflections $r_0, \dots, r_{N-1}$. If $1 \leq \sigma \leq N-1$, $r_\sigma := \mathcal{E}_\sigma \leftrightarrow \mathcal{E}_{\sigma+1}$ is the reflection about the element $\mathcal{E}_\sigma - \mathcal{E}_{\sigma+1}$; and r_0 is the Cremona involution, which is the reflection about $\mathcal{E}_0 - \mathcal{E}_1 - \mathcal{E}_2 - \mathcal{E}_3$. The Coxeter is the product $r_1 \cdots r_{N-1} r_0$ of these involutions. Writing the product $r_1 \cdots r_{N-1}$ in a different order corresponds to permuting the basis elements $\mathcal{E}_\sigma$, $1 \leq \sigma \leq N$. Thus we may write the Coxeter element (after a possible permutation of basis elements) as:

$$\begin{aligned} \mathcal{E}_0 &\mapsto 2\mathcal{E}_0 - \mathcal{E}_2 - \mathcal{E}_3 - \mathcal{E}_4 \\ \mathcal{E}_1 &\mapsto \mathcal{E}_0 - \mathcal{E}_3 - \mathcal{E}_4 \\ \mathcal{E}_2 &\mapsto \mathcal{E}_0 - \mathcal{E}_2 - \mathcal{E}_4 \\ \mathcal{E}_3 &\mapsto \mathcal{E}_0 - \mathcal{E}_2 - \mathcal{E}_3 \\ \mathcal{E}_\sigma &\mapsto \mathcal{E}_{\sigma+1}, \quad 4 \leq \sigma \leq N-1, \quad \mathcal{E}_N \mapsto \mathcal{E}_1. \end{aligned}$$

Theorem 8.1. Suppose that $\mathcal{X}$ is a rational surface, $F \in \text{Aut}(\mathcal{X})$ represents the Coxeter element of W_N for $N \geq 5$, and suppose that we represent $\mathcal{X}$ by iterated blowups $\pi : \mathcal{X} \rightarrow$

$\mathbf{P}^2$. Then there exist a linear map $T \in \text{Aut}(\mathbf{P}^2)$ and complex numbers a and b such that

$$f_{a,b} \circ T \circ \pi = T \circ \pi \circ F.$$

Before giving the proof, let us recall that for $\sigma \geq 1$, the $\mathcal{E}_\sigma$ are given as $E_\sigma + \sum_t \mu_{\sigma,t} E_t$, where $\mu_{\sigma,t}$ is a nonnegative integer, and E_t is an irreducible exceptional fiber “over” E_σ . In addition, each of the basis elements $\mathcal{E}_\sigma$, $1 \leq \sigma \leq N$ lies over a single point $p_\sigma := \pi(E_\sigma)$. It is uniquely represented in terms of prime components, so the number $\#\mathcal{E}_\sigma$ of these prime components is well defined for $1 \leq \sigma \leq N$.

Lemma 8.2. *We have $\#\mathcal{E}_\sigma = 1$, i.e., $\mathcal{E}_\sigma = E_\sigma$ for $4 \leq \sigma \leq N$, and $\sigma = 1$.*

Proof. Since the representation $E_\sigma + \sum_t \mu_{\sigma,t} E_t$ is unique, we have $\#\mathcal{E}_4 = \dots = \#\mathcal{E}_N = \#\mathcal{E}_1$. Now if $\#\mathcal{E}_4 > 1$, then there is some $\tau = \tau_4$ such that $\mathcal{E}_{\tau_4}$ lies strictly above $\mathcal{E}_4$, and thus $\#\mathcal{E}_4 > \#\mathcal{E}_{\tau_4} \geq 1$. Now F_* maps $\mathcal{E}_{\tau_4}$ to some basis element $\mathcal{E}_{\tau_5}$, which lies strictly above $\mathcal{E}_5$. It is evident that we have $\#\mathcal{E}_{\tau_4} = \#\mathcal{E}_{\tau_5} = \#\mathcal{E}_{\tau_1} < \#\mathcal{E}_4$, so we have three new independent elements. But now we have too many independent elements for the dimension of $\text{Pic}(\mathcal{X})$, so we conclude that we must have had $\#\mathcal{E}_4 = 1$. $\square$

Lemma 8.3. *The points p_2 and p_3 are distinct.*

Proof. The Coxeter element above maps $\mathcal{E}_3 \mapsto \mathcal{E}_0 - \mathcal{E}_2 - \mathcal{E}_3 \mapsto \mathcal{E}_2$, so $\#\mathcal{E}_2 = \#\mathcal{E}_3$. If $\#\mathcal{E}_3 = \#\mathcal{E}_2 = 1$, then we have $\#\mathcal{E}_\sigma = 1$ for all $1 \leq \sigma \leq N$. Thus the points p_σ are distinct. Now suppose that $\#\mathcal{E}_3 = \#\mathcal{E}_2 > 1$. Then in the representation $\mathcal{E}_3 = E_3 + \sum_t \mu_{3,t} E_t$ there are components E_t lying above E_3 . However, if $p_2 = p_3$, then either E_2 lies above E_3 or E_3 lies above E_2 . In the first case, for instance, we see from this representation that we must have $\#\mathcal{E}_2 < \#\mathcal{E}_3$. Thus we must have $p_2 \neq p_3$. $\square$

Lemma 8.4. *The points p_1, p_2, p_3 and p_4 are distinct*

Proof. Let us start by showing that various of the p_i ’s are distinct. First we show $p_1 \neq p_2$. Suppose not. If $p_1 = p_2$, then since $\#\mathcal{E}_1 = 1$, we have $\mathcal{E}_1 = E_1$, and the fiber E_1 must lie above E_2 . And in the representation $\mathcal{E}_2 = E_2 + E_1 + \sum_{t \neq 1,2,3} \mu_{2,t} E_t$, and all the other terms $\mu_{2,t}$ are either 0 or 1. Thus

$$E_2 = \mathcal{E}_2 - \mathcal{E}_1 - \sum_{t \neq 1,2,3} \mu_{2,t} E_t,$$

and mapping this expression forward by the Coxeter element gives

$$E_2 \mapsto \mathcal{E}_3 - \mathcal{E}_2 - \sum_{t \neq 1,2,3} \mu_{2,t} E_{t+1}.$$

But this is not possible, since E_2 is positive and irreducible, but the image in the line above is neither positive nor irreducible, since $\mathcal{E}_2$ and $\mathcal{E}_3$ are carried over distinct points $p_2 \neq p_3$.

Similarly, we show that $p_2 \neq p_4$. For if $p_2 = p_4$, then we must have E_4 above E_2 , and in the expression $\mathcal{E}_2 = E_2 + \sum_t \mu_{2,t} E_t$, we have $\mu_{2,4} = 1$. The inverse of the Coxeter element acts like $\mathcal{E}_2 \mapsto \mathcal{E}_0 - \mathcal{E}_2 - \mathcal{E}_3$ and $\mathcal{E}_4 \mapsto \mathcal{E}_0 - \mathcal{E}_1 - \mathcal{E}_2$. Thus the inverse acts by

$$E_2 \mapsto \mathcal{E}_1 - \mathcal{E}_3 - \sum_{t \neq 2,3,4} \mu_{2,t} E_{t-1},$$

which is not possible since the right hand side is not positive since $\mathcal{E}_3 - \mathcal{E}_1 \geq E_3$.

Next we show that $p_1 \neq p_3$ and $p_4 \neq p_3$ by the same argument. Finally, we see that $p_1 \neq p_4$. Since p_1 and p_4 are disjoint from $\{p_2, p_3\}$, it follows that there is just one level of blowup in $\mathcal{X}$ over p_1 and p_4 . Since $E_\sigma = \mathcal{E}_\sigma$ for $\sigma = 1, 4$, and these are distinct elements of $\text{Pic}(\mathcal{X})$, $p_1 \neq p_4$. $\square$

Lemma 8.5. $p_1, p_4 \notin \overline{p_2 p_3}$.

Proof. By Lemma 8.2 the complete fiber over p_1 is $E_1 = \mathcal{E}_1$, which is irreducible. Thus the image $F(E_1)$ is an irreducible curve in $\mathcal{X}$ which represents $\mathcal{E}_0 - \mathcal{E}_3 - \mathcal{E}_4$. The curve $\pi(F(E_1))$ is a curve in $\mathbf{P}^2$ of degree 1 and thus is a line which must pass through p_3 and p_4 . Since $F(E_1)$ is irreducible, it cannot pass through p_1 or p_2 . In particular, the points p_2, p_3, p_4 are not colinear, so $p_4 \notin \overline{p_2 p_3}$. Similarly, we apply the inverse of the Coxeter element to $E_4 = \mathcal{E}_4$ and conclude that p_3 and p_4 are not on the line through $p_1 p_2$. Thus p_1, p_2 and p_3 are not collinear, so $p_1 \notin \overline{p_2 p_3}$. $\square$

Proof of Theorem 8.1. After Lemmas 8.2–5, we are in the situation of Theorem 11.1 of [M2]. We give the proof here for the sake of completeness. Let φ be the birational map of $\mathbf{P}^2$ induced by F . Since $p_1 \notin \overline{p_2 p_3}$, the points p_1, p_2 and p_3 are the vertices of a proper triangle. Thus we may introduce a linear change of coordinates $[t : x : y]$ such that $p_1 = [1 : 0 : 0]$, $p_2 = [0 : 1 : 0]$, and $p_3 = [0 : 0 : 1]$. Now define α and β by the condition that $p_4 = [1 : \alpha : \beta]$. Define $g_{\alpha,\beta} = (y, y/x) + (\alpha, \beta)$. Then since φ and $g_{\alpha,\beta}$ have the same exceptional curves and map them in the same way, there are constants A and B such that $\varphi(Ax, By) = g_{\alpha,\beta}(x, y)$. Since $g_{\alpha,\beta}$ is linearly conjugate to some $f_{a,b}$, the Theorem follows. $\square$

Let $\pi : \mathcal{Y} \rightarrow \mathbf{P}^2$ denote the space obtained by blowing up the points e_1 and e_2 in $\mathbf{P}^2$. Every map $f_{a,b}$ induces a birational map $f_{\mathcal{Y}}$ of $\mathcal{Y}$ to itself, and $p = (-b, -a)$ is the only point of indeterminacy for $f_{\mathcal{Y}}$. Thus the pointwise iteration $f_{\mathcal{Y}}^j := f_{\mathcal{Y}} \circ \cdots \circ f_{\mathcal{Y}}$ is well defined until the orbit reaches p . In other words:

Lemma 8.6. *If $r \in \mathcal{Y}$ and $f_{\mathcal{Y}}^j r \neq p$ for $0 \leq j \leq k$, then $(f^k)_{\mathcal{Y}}(r) = f_{\mathcal{Y}} \circ \cdots \circ f_{\mathcal{Y}}(r)$, there the composition on the right is k -fold.*

The point of Lemma 8.6 is that since we may identify $\mathbf{C}^2$ with $\mathcal{Y} - (E_1 \cup E_2 \cup \Sigma_0)$, we may define $\mathcal{V}_n$ in terms of $f_{\mathcal{Y}}$:

$$\mathcal{V}_n = \{(a, b) \in \mathbf{C}^2 : f_{\mathcal{Y}}^j q \neq p \text{ for } 0 \leq j < n, f_{\mathcal{Y}}^n q = p\}.$$

By Lemma 8.2, a representation of the Coxeter element can be obtained by making simple (not iterated) blowups of birational maps $f_{a,b}$ which have been lifted to $\mathcal{Y}$. We state it as follows:

Theorem 8.7. Suppose that $\mathcal{X}$ is a rational surface, and $F \in \text{Aut}(\mathcal{X})$ represents the Coxeter element in W_N . Then there exists a surface $\pi : \mathcal{Z} \rightarrow \mathcal{Y}$ which is obtained by the blowup of a finite set of distinct points in $\mathcal{Y}$, and there are an automorphism $G \in \text{Aut}(\mathcal{Z})$ and $(a, b) \in \mathcal{V}_{N-3}$ such that $(F, \mathcal{X})$ is conjugate to $(G, \mathcal{Z})$, and $\pi \circ G = f_{\mathcal{Y}} \circ \pi$, with $f_{\mathcal{Y}} = (f_{a,b})_{\mathcal{Y}}$.

Remarks. Theorems 8.1 and 8.7 give a strengthening of Theorem 11.1 of [M2] so that it applies to all rational surface maps. We note some notational differences with [M2]. Namely, [M2] uses N for the number of blowups, whereas [BK] and the present paper use n for the length of the orbit of q . These two numbers are related by $N = n + 3$. The 2-dimensional surface on which an automorphism is defined is denoted by $\mathcal{X}_{a,b}$ in [BK], whereas the surface is S in [M2], and X represents an invariant curve. Here we use the notation (0.1) of [BK] for the birational family; [M2] changed this notation to $(x, y) \mapsto (y, y/x) + (\alpha, \beta)$, which is conjugate to $f_{a,b}$ if $\alpha = b - a$ and $\beta = a$.

Appendix A. Varieties $\mathcal{V}_j$ and Γ for $0 \leq j \leq 6$. The sets $\mathcal{V}_j$, $0 \leq j \leq 6$ are enumerated in [BK]. We note that $\mathcal{V}_0 = (0, 0) \subset \Gamma_2 \cap \Gamma_3$, $\mathcal{V}_1 = (1, 0) \subset \Gamma_1$, $\mathcal{V}_2 \subset \Gamma_1 \cap \Gamma_2$, $\mathcal{V}_3 \subset \Gamma_1 \cap \Gamma_3$, and $(\mathcal{V}_4 \cup \mathcal{V}_5) \cap \Gamma = \emptyset$.

Each of the mappings in $\mathcal{V}_6$ has an invariant pencil of cubics. There are two cases: the set $\mathcal{V}_6 \cap \{b \neq 0\}$ (consisting of four points) is contained in $\Gamma_2 \cap \Gamma_3$. The other case, $\mathcal{V}_6 \cap \{b = 0\} = \{(a, 0) : a \neq 0, 1\}$, differs from the cases $\mathcal{V}_n$, $n \neq 0, 1, 6$, because the manifold $\mathcal{X}_{a,0}$ is constructed by iterated blowups ($f^4 q \in E_1$ and $f^2 q \in E_2$, see [BK, Figure 6.2]).

The invariant function $r(x, y) = (x + y + a)(x + 1)(y + 1)/(xy)$ for $f_{a,0}$, which defines the invariant pencil, was found by Lyness [L] (see also [KLR], [KL], [BC] and [Z1]). We briefly describe the behavior of $f_{a,0}$. By $M_\kappa = \{r = \kappa\}$ we denote the level set of r inside $\mathcal{X}_{a,0}$. The curve M_∞ consists of an invariant 5-cycle of curves with self-intersection -2 :

$$\Sigma_\beta = \{x = 0\} \mapsto E_2 \mapsto \Sigma_0 \mapsto E_1 \mapsto \Sigma_B = \{y = 0\} \mapsto \Sigma_\beta.$$

The restriction of f^5 to any of these curves is a linear (fractional) transformation, with multipliers $\{a, a^{-1}\}$ at the fixed points. M_0 consists of a 3-cycle of curves with self-intersection -1 : $\{y + 1 = 0\} \mapsto \{x + 1 = 0\} \mapsto \{x + y + a = 0\}$. The restriction of f^3 to any of these lines is linear (fractional) with multipliers $\{a - 1, (a - 1)^{-1}\}$ at the fixed points.

Theorem A.1. Suppose that $a \notin \{-\frac{1}{4}, 0, \frac{3}{4}, 1, 2\}$, and $\kappa \neq 0, \infty$. If M_κ contains no fixed point, then M_κ is a nonsingular elliptic curve, and f acts as translation on M_κ . If M_κ contains a fixed point p , then M_κ has a node at p . If we uniformize $s : \hat{\mathbf{C}} \rightarrow M_\kappa$ so that $s(0) = s(\infty) = p$, then $f|_{M_\kappa}$ is conjugate to $\zeta \mapsto \alpha\zeta$ for some $\alpha \in \mathbf{C}^*$.

The intersection $\Gamma_j \cap \{b = 0\} \cap \mathcal{V}_6$ is given by $(-\frac{1}{4}, 0)$, $(\frac{3}{4}, 0)$, or $(0, 2)$, if $j = 1, 2$, or 3 , respectively.

Theorem A.2. Suppose that $a = -\frac{1}{4}, \frac{3}{4}$, or 2 . Then the conclusions of Theorem A.1 hold, with the following exception. If $FP_s \in M_\kappa$, then M_κ is a cubic which has a cusp at FP_s , or is a line and a quadratic tangent at FP_s , or consists of three lines passing through

FP_s . If we uniformize a component $s : \hat{\mathbf{C}} \rightarrow M_\kappa$ such that $s(\infty) = FP_s$, then $f^j|_{M_\kappa}$ is conjugate to $\zeta \mapsto \zeta + 1$, where j is chosen so that $(a, 0) \in \Gamma_j$.

Appendix B. Varieties $\mathcal{V}_n$ and Γ for $n \geq 7$. We may compute $\mathcal{V}_n$ explicitly by starting with the equation $f_{a,b}^n(-a, 0) + (b, a) = 0$. We look upon $f_{a,b}^n(-a, 0)$ as a rational expression in a and b with integer coefficients, so we first remove common factors from the numerator and denominator of f^n , and then we convert this equation to a pair of polynomial equations $S_n = \{P_n(a, b) = Q_n(a, b) = 0\}$ with integer coefficients in the variables a and b . Thus S_n contains all the elements of $\mathcal{V}_n$ and possibly more. We may find the elements of S_n by applying elimination theory to P_n and Q_n ; since they are polynomials with integer coefficients, we obtain an exact polynomial for the possible values of a (or b). We may then test numerically whether a given element $(a, b) \in S_n$ actually belongs to $\mathcal{V}_n$, and not some $\mathcal{V}_j$ for $j < n$, by computing the orbit $f_{a,b}^j q$, $1 \leq j \leq n$. We do this using *Mathematica* (or *Maple*) and find that $\#\mathcal{V}_7 = 10$. On the other hand, by Theorem 3.5, $\chi_7(x) = (x-1)\psi_7(x)$, and so ψ_7 has degree 10. By Theorem 3.4, $\#(\mathcal{V}_7 \cap \Gamma_1) = 10$, and so by counting we conclude that $\mathcal{V}_7 \subset \Gamma_1$. Arguing in this manner, we obtain

Proposition B.1. $\mathcal{V}_7 \subset \Gamma_1$, $\mathcal{V}_8 \cup \mathcal{V}_{10} \subset \Gamma_1 \cup \Gamma_2$, and $\mathcal{V}_9 \subset \Gamma_1 \cup \Gamma_3$.

Counterexamples. To find the counterexamples discussed in §4, we let $n = 11$. By Theorem 3.5, ψ_{11} has degree 10, so by Theorem 3.3, $\Gamma_1 \cap \mathcal{V}_{11} = \Gamma \cap \mathcal{V}_{11}$ contains 10 elements. On the other hand, we may use *Mathematica* to determine the number of elements of $\mathcal{V}_{11}$. That is, we follow the scheme described in the previous paragraph and obtain the resultant of P_{11} and Q_{11} , eliminating either a or b . This resultant has a factor of degree 10 and one of degree 12. The 10 roots of the degree 10 factor correspond to the values of $\Gamma \cap \mathcal{V}_{11}$, and the 12 roots of the degree 12 factor correspond to the values of $\mathcal{V}_{11} - \Gamma$, and thus these 12 maps do not have invariant curves.

We have repeated this computational procedure in the cases $11 < n \leq 25$, and we have found that $\mathcal{V}_n \cap \Gamma$ becomes a quite small fraction of $\mathcal{V}_n$ as n increases.

A map $f_{a,b}$ has a unique 2-cycle and a unique 3-cycle. For $\ell = 2, 3$, we let J_ℓ denote the product of the Jacobian matrix around the ℓ -cycle.

Theorem B.2. For $\ell = 2, 3$, the determinant μ_ℓ of J_ℓ is given by

$$\mu_2 = \frac{a - b - 1}{2b^2 + a - 1}, \quad \mu_3 = \frac{1 + b + b^2 - a - ab}{1 - a - ab} \quad (B.1)$$

and the trace τ_ℓ is given by

$$\tau_2 = \frac{3 - 2a + b - b^2}{2b^2 + a - 1}, \quad \tau_3 = \frac{2 + a^2 + b + 2b^2 - b^3 + b^4 + a(-2 - b + 2b^2)}{-1 + a - ab}. \quad (B.2)$$

Proof. The proof of the 2-cycle case is simpler and omitted. We may identify a 3-cycle with a triple of numbers z_1, z_2, z_3 :

$$\zeta_1 = (z_1, z_2) \mapsto \zeta_2 = (z_2, z_3 = \frac{a + z_2}{b + z_1}) \mapsto \zeta_3 = (z_3, z_1 = \frac{a + z_3}{b + z_2}) \mapsto \zeta_1 = (z_1, z_2 = \frac{a + z_1}{b + z_3}).$$

Substituting into this 3-cycle, we find that z_1, z_2, z_3 are the three roots of

$$P_3(z) = z^3 + (1 + a + b + b^2)z^2 + (b^3 + ab + 2a - 1)z - 1 + a - b + ab - b^2.$$

It follows that

$$\begin{aligned} z_1 + z_2 + z_3 &= -(1 + a + b + b^2) \\ z_1 z_2 + z_1 z_3 + z_2 z_3 &= -1 + 2a + ab + b^3 \\ z_1 z_2 z_3 &= 1 - a + b - ab + b^2 \end{aligned} \tag{B.3}$$

Since

$$Df_{a,b}(\zeta_1 = (z_1, z_2)) = \begin{pmatrix} 0 & 1 \\ -\frac{a+z_2}{(b+z_1)^2} & \frac{1}{b+z_1} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ \frac{-z_3}{b+z_1} & \frac{1}{b+z_1} \end{pmatrix},$$

the determinant of $Df_{a,b}(\zeta_1) = z_3/(b + z_1)$, and therefore

$$\mu_3 = \frac{z_2}{b + z_3} \frac{z_1}{b + z_2} \frac{z_3}{b + z_1} = \frac{z_1 z_2 z_3}{(b + z_3)(b + z_2)(b + z_1)}.$$

Using equations (B.3) we see that $(b + z_1)(b + z_2)(b + z_3) = 1 - a + ab$ so μ_3 has the form given in (B.1).

Similarly, we compute

$$Tr(J_3) = -\frac{-1 + b(z_1 + z_2 + z_3) + z_1^2 + z_2^2 + z_3^2}{(b + z_1)(b + z_2)(b + z_3)}.$$

Using (B.3) again, we find that τ_3 is given in the form (B.2). □

A computation shows the following:

Theorem B.3. *Suppose that $(a, b) = \varphi_1(t) \in \mathcal{V}_n \cap \Gamma_1$, $n \geq 7$. Then the eigenvalues of Df at FP_r are given by $\{\eta_1 = 1/t, \eta_2 = -(t^3 + t^2 - 1)/(t^4 - t^2 - t)\}$, where t is a root of ψ_n . Further, they have the resonance $\eta_1^n \eta_2 = 1$.*

In the previous theorem, the fixed point FP_r is contained in the invariant curve. If ℓ divides n , for $\ell = 2$ or 3 , then the ℓ -cycle is disjoint from the invariant curve. Thus we have:

Theorem B.4. *Suppose $\ell = 2$ or 3 and $n = k\ell \geq 7$. If $(a, b) = \varphi_\ell(t) \in \mathcal{V}_n \cap \Gamma_\ell$, then the eigenvalues of the ℓ -cycle are $\{\eta_1 = t^{-\ell}, \eta_2 = -t^{\ell-1}(t^3 + t^2 - 1)/(t^3 - t - 1)\}$. Further, they have the resonance $\eta_1^{n+1} \eta_2 = 1$.*

Corollary B.5. *If $t = \lambda_n$ or λ_n^{-1} , then the cycles discussed in Theorems B.3 and B.4 are saddles. If t has modulus 1, then the multipliers over these cycles have modulus 1 but are not roots of unity.*

Appendix C. Computation of Characteristic Polynomials.

Theorem C.1. *If χ_n is as in (0.3), and $n \geq 7$, then*

- (i) *The characteristic polynomial for (6.1) is $(x^7 + 1)\chi_n(x)/(x^2 - 1)$;*
- (ii) *The characteristic polynomial for (6.2) is $(x^5 - 1)\chi_{2k}(x)/(x^2 - 1)$;*
- (iii) *The characteristic polynomial for (6.3) is $(x^4 - 1)\chi_{3k}(x)/(x^3 - 1)$.*

Proof. We start with case (i). Since the case $n = 7$ is easily checked directly, it suffices to prove (i) for $n \geq 8$. Let us use the ordered basis:

$$\{12, 23, 34, 04, 15, 26, 03, 14, 25, 05, 16, 02, 13, 24, 06, 170, 180, \dots, 1n0\},$$

and let $M = (m_{i,j})$ denote the matrix which represents the transformation $\eta \mapsto f_*\eta$ defined in (5.1), i.e., we set $m_{i,j} = 1$ if the i -th basis element in our ordered basis maps to the j -th basis element, and 0 otherwise. To compute the characteristic polynomial of M , we expand $\det(M - xI)$ by minors down the last column. We obtain

$$\det(M - xI) = -xM_{n+9,n+9} + (-1)^n M_{1,n+9}, \quad (C.1)$$

where we use the notation $M_{i,j}$ for the i, j -minor of the matrix $M - xI$. To evaluate $M_{n+9,n+9}$ and $M_{1,n+9}$, we expand again in minors along the last column to obtain

$$M_{n+9,n+9} = -x \det \hat{m}_1 + (-1)^n \det \hat{m}_2, \quad M_{1,n+9} = \det \hat{m}_3,$$

where $\hat{m}_1 = \begin{pmatrix} A_1 & 0 \\ 0 & A_2(n) \end{pmatrix}$, $\hat{m}_2 = \begin{pmatrix} B_1 & * \\ 0 & B_2(n) \end{pmatrix}$, and $\hat{m}_3 = \begin{pmatrix} C_1 & * \\ 0 & C_2(n) \end{pmatrix}$. Here A_1 , B_1 , and C_1 do not depend on n , and $A_2(n)$, $B_2(n)$, and $C_2(n)$ are triangular matrices of size $(n-7) \times (n-7)$, $(n-8) \times (n-8)$ and $(n-8) \times (n-8)$ of the form

$$A_2(n) = \begin{pmatrix} -x & 0 & 0 \\ * & \ddots & 0 \\ * & * & -x \end{pmatrix}, \quad B_2(n) = \begin{pmatrix} 1+x & * & * \\ 0 & \ddots & * \\ 0 & 0 & 1+x \end{pmatrix}, \quad C_2(n) = \begin{pmatrix} 1 & * & * \\ 0 & \ddots & * \\ 0 & 0 & 1 \end{pmatrix}.$$

Thus

$$\det A_2(n) = (-x)^{n-7}, \quad \det B_2(n) = (1+x)^{n-8}, \quad \text{and} \quad \det C_2(n) = 1. \quad (C.2)$$

Since A_1 , B_1 , and C_1 do not depend on n , we may compute them using the matrix M from the case $n = 8$ to find

$$\det A_1 = -x^6(x^8 - x^5 - x^3 + 1), \quad \det B_1 = -x^8, \quad \text{and} \quad \det C_1 = x^5 + x^3 - 1. \quad (C.3)$$

Using (C.2) and (C.3) we find that the characteristic polynomial of M is equal to

$$(-1)^n [x^9(x+1)^{n-8} - x^{n+1}(x^8 - x^5 - x^3 + 1) + x^5 + x^3 - 1] = (x^7 - 1)\chi_n(x)/(x^2 - 1),$$

which completes the proof of (i).

For the proof of (ii), we use the ordered basis

$$\{12, 2a3, 34, 45, 1g2, 2e3, 3c4, 04, 01, 03, 14, 25, 061, 072, 081, \dots, 0(2k-1)2, 0(2k)1\},$$

and for (iii) we use the ordered basis

$$\{23, 3a4, 015, 13, 01, 12, 2d3, 3c4, 02, 04, 162, 073, 081, 192, \dots, 0(3k-1)1, 1(3k)2\}.$$

Otherwise, the proofs of cases (ii) and (iii) are similar. We omit the details. $\square$

References.

- [B] A. Baker, The theory of linear forms in logarithms, in *Transcendence Theory: Advances and Applications*, edited by A. Baker and D.W. Masser, Academic Press 1977, pp. 1–27.
- [BD1] E. Bedford and J. Diller, Real and complex dynamics of a family of birational maps of the plane: the golden mean subshift. *Amer. J. Math.* 127 (2005), no. 3, 595–646.
- [BD2] E. Bedford and J. Diller, Dynamics of a two parameter family of plane birational maps: maximal entropy, *J. of Geometric Analysis*, 16 (2006), no. 3, 409–430.
- [BK] E. Bedford and KH Kim, Periodicities in Linear Fractional Recurrences: Degree growth of birational surface maps, *Mich. Math. J.*, 54 (2006), 647–670.
- [BLS] E. Bedford, M. Lyubich, and J. Smillie, Polynomial diffeomorphisms of C^2 . IV. The measure of maximal entropy and laminar currents. *Invent. Math.* 112 (1993), no. 1, 77–125.
- [BS1] E. Bedford and J. Smillie, Polynomial diffeomorphisms of C^2 . II. Stable manifolds and recurrence. *J. Amer. Math. Soc.* 4 (1991), no. 4, 657–679.
- [BS2] E. Bedford and J. Smillie, Real polynomial diffeomorphisms with maximal entropy: Tangencies. *Ann. of Math.* (2) 160 (2004), no. 1, 1–26.
- [BC] F. Beukers and R. Cushman, Zeeman’s monotonicity conjecture, *J. of Differential Equations*, 143 (1998), 191–200.
- [C1] S. Cantat, Dynamique des automorphismes des surfaces projectives complexes. *C. R. Acad. Sci. Paris Sér. I Math.* 328 (1999), no. 10, 901–906.
- [C2] S. Cantat, Dynamique des automorphismes des surfaces $K3$. *Acta Math.* 187 (2001), no. 1, 1–57.
- [DJS] J. Diller, D. Jackson, and A. Sommesse, Invariant curves for birational surface maps, *Trans. AMS* 359 (2007), 2973–2991. [math.AG/0505014](#)
- [F] W. Floyd, Growth of planar Coxeter groups, P.V. numbers, and Salem numbers, *Math. Ann.* 293, 475–483 (1992).
- [FS] J.-E. Fornæss and N. Sibony, Classification of recurrent domains for some holomorphic maps. *Math. Ann.* 301 (1995), no. 4, 813–820.
- [G] M. Gromov, Entropy, homology and semialgebraic geometry. *Séminaire Bourbaki*, Vol. 1985/86. *Astérisque* No. 145-146 (1987), 5, 225–240.
- [H] B. Harbourne, Rational surfaces with infinite automorphism group and no antipluri-canonical curve, *Proc. AMS*, 99 (1987), 409–414.
- [KLR] V.I. Kocic, G. Ladas, and I.W. Rodrigues, On rational recursive sequences, *J. Math. Anal. Appl* 173 (1993), 127–157.
- [KuL] M. Kulenovic and G. Ladas, *Dynamics of Second Order Rational Difference Equations*, CRC Press, 2002.
- [L] R.C. Lyness, Notes 1581, 1847, and 2952, *Math. Gazette* **26** (1942), 62, **29** (1945), 231, and **45** (1961), 201.
- [M1] C. McMullen, Dynamics on $K3$ surfaces: Salem numbers and Siegel disks. *J. Reine Angew. Math.* 545 (2002), 201–233.
- [M2] C. McMullen, Dynamics on blowups of the projective plane, *Pub. Sci. IHES*, 105 (2007), p. 49–89.

- [P] J. Pöschel, On invariant manifolds of complex analytic mappings near fixed points. *Exposition. Math.* 4 (1986), No. 2, 97–109.
- [U] T. Ueda, Critical orbits of holomorphic maps on projective spaces, *J. of Geometric Analysis*, 8 (1998), 319–334.
- [W] L. Washington, *Introduction to Cyclotomic Fields*, Springer-Verlag 1997.
- [Y] Y. Yomdin, Volume growth and entropy. *Israel J. Math.* 57 (1987), no. 3, 285–300.
- [Z1] C. Zeeman, Geometric unfolding of a difference equation, lecture.
- [Z2] E. Zehnder, A simple proof of a generalization of a theorem by C. L. Siegel. *Geometry and topology* (Proc. III Latin Amer. School of Math., Inst. Mat. Pura Aplicada CNPq, Rio de Janeiro, 1976), pp. 855–866. *Lecture Notes in Math.*, Vol. 597, Springer, Berlin, 1977.
- [Zh] D-Q Zhang, Automorphism groups and anti-canonical curves, *Math. Res. Lett.* 15 (2008), 163–183.

Indiana University
Bloomington, IN 47405
bedford@indiana.edu

Florida State University
Tallahassee, FL 32306
kim@math.fsu.edu